

HIV Surveillance **Report** | Supplemental Report

Volume 21, Number 2

Diagnoses of HIV Infection among Adults Aged 50 Years and Older in the United States and Dependent Areas 2010–2014

This issue of the *HIV Surveillance Supplemental Report* is published by the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, Georgia.

Data are presented for diagnoses of HIV infection reported to CDC through July 2015.

The *HIV Surveillance Supplemental Report* is not copyrighted and may be used and copied without permission. Citation of the source is, however, appreciated.

Suggested citation

Centers for Disease Control and Prevention. Diagnoses of HIV infection among adults aged 50 years and older in the United States and dependent areas, 2010–2014. *HIV Surveillance Supplemental Report* 2016;21(No. 2). <http://www.cdc.gov/hiv/library/reports/surveillance/>. Published May 2016. Accessed [date].

On the Web: <http://www.cdc.gov/hiv/library/reports/surveillance/>.

Confidential information, referrals, and educational material on HIV infection

CDC-INFO

1-800-232-4636 (in English, en Español)

1-888-232-6348 (TTY)

<http://www.cdc.gov/cdc-info/requestform.html>

Acknowledgments

Publication of this report was made possible with the contributions of the state and territorial health departments and the HIV surveillance programs that provided surveillance data to CDC.

This report was prepared by the following staff and contractors of the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, CDC:

Benjamin Laffoon, Laurie Linley, Jing Shi, Jianmin Li, H. Irene Hall, Anna Satcher Johnson, and Michael Friend (desktop publishing).

Contents

Commentary	5
Technical Notes	9
References	13
Tables	
Section 1 Diagnoses of HIV Infection among Adults Aged 50 Years and Older	
1a Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States	14
1b Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States and 6 dependent areas	16
2a Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States	18
2b Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States and 6 dependent areas	20
3a Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States	22
3b Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States and 6 dependent areas	24
4a Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States	26
4b Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States and 6 dependent areas	28
5 Diagnoses of HIV infection among adults aged 50 years and older, by age group and area of residence, 2014—United States and 6 dependent areas	30
Section 2 Adults Aged 50 Years and Older Living with Diagnosed HIV Infection (Prevalence)	
6a Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States	32
6b Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas	34
7a Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States	36
7b Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States and 6 dependent areas	38
8a Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States	40

8b	Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States and 6 dependent areas	42
9	Adults aged 50 years and older living with diagnosed HIV infection, by age group and area of residence, year-end 2013—United States and 6 dependent areas	44
10	Adults aged 50 years and older living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas	46
11	Adults aged 50 years and older living with diagnosed HIV infection, by population of area of residence, age group, and selected characteristics, year-end 2013—United States	48

Section 3 Deaths of Adults Aged 50 Years and Older with Diagnosed HIV Infection

12a	Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States	50
12b	Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas	52
13a	Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States	54
13b	Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States and 6 dependent areas	56
14a	Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States	58
14b	Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States and 6 dependent areas	60

Section 4 Stage 3 (AIDS) Classification at the Time of Diagnosis of HIV Infection among Adults Aged 50 Years and Older

15a	Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States	62
15b	Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas	64
16a	Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by age group and selected characteristics, 2013—United States	66
16b	Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas	68

Data from the 2014 *HIV Surveillance Report*, published by the Centers for Disease Control and Prevention (CDC), indicate that the estimated percentage of persons aged 50 years and older and living with diagnosed HIV infection increased from 35% in 2010 to 42% in 2014 [1].

This surveillance supplemental report complements the 2014 *HIV Surveillance Report* by presenting data on diagnoses of HIV infection among persons aged 50 years and older in the United States and 6 dependent areas. In this report, the term *older* refers to adults aged 50 years and older. This report is intended to guide public health strategies that focus on preventing the transmission of HIV among older adults in the United States and dependent areas as well as to assist with planning for HIV treatment and care needs.

The term *diagnosis of HIV infection* refers to a diagnosis of HIV infection regardless of the stage of disease (stage 1, 2, 3 [AIDS], or unknown) and duration of infection at the time of diagnosis. In this report, the 2008 and 2014 HIV case definitions were used to classify HIV infection among adults aged 50 years and older. The 2008 case definition was used to classify cases diagnosed from 2010 through 2013. The 2014 case definition was used to classify cases diagnosed in 2014. For more information on the stages of HIV infection, please review the Technical Notes.

REPORT FORMAT

This report presents the unadjusted numbers and the estimated (statistically adjusted for reporting delays) numbers and rates (cases diagnosed per 100,000 persons) of HIV diagnosed infection during 2010–2014 for older adults. Data are also presented for older adults living with diagnosed HIV infection (prevalence) and deaths of older adults with diagnosed HIV infection as of December 31, 2013. Deaths of persons with diagnosed HIV infection may be due to any cause (i.e., may or may not be related to HIV infection). Data are also presented for older adults whose HIV infection was diagnosed during 2013 and was classified as stage 3 (AIDS) at the time of HIV diagnosis.

HIGHLIGHTS OF ANALYSES

The statements in this section, unless otherwise indicated, are based on estimated (statistically adjusted) data. All rates are per 100,000 population. The standard used for reporting trends in numbers and rates is based on an increase or a decrease of 5% or more during the specified time frame (e.g., when comparing 2010 and 2014). Trends based on fewer than 12 diagnoses or rates based on small numbers were not included in the narrative of this report. Please read all table titles and footnotes carefully to ensure a complete understanding of the displayed data.

Diagnoses of HIV Infection among Adults Aged 50 Years and Older

From 2010 through 2014, the estimated number of annual diagnoses of HIV infection among older adults in the United States remained stable, but the rate decreased. In 2014, the estimated rate was 6.8 per 100,000 population (Table 1a).

- **Age group:** From 2010 through 2014, the rates for adults aged 50–54, 55–59, 60–64, 65–69, and 70–74 years decreased. The rates remained stable for adults aged 75–79 and 80–84 years. In 2014, the largest percentage of all diagnoses (44%) and the highest rate per 100,000 population (14.4) were those for adults aged 50–54 years (Table 1a). The estimated rate of diagnosed HIV infections for adults aged 50–64 years (10.3) was 5 times the rate for adults aged 65 years and older (2.0) (Table 2a). Please use caution when interpreting data for adults aged 85 years and older: the numbers are small.
- **Race/ethnicity:** From 2010 through 2014, the rates for Asians increased. The rates for American Indians/Alaska Natives, blacks/African Americans, Hispanics/Latinos, and adults of multiple races decreased. The rates for whites remained stable (Table 1a). In 2014, blacks/African Americans accounted for 43% of all diagnoses among older adults. The rate for blacks/African Americans (28.4) was 2 times the rate for Hispanics/Latinos (11.4) and 8 times the rate for whites (3.4) (Table 1a). Please use cau-

tion when interpreting data for Native Hawaiians/ other Pacific Islanders: the numbers are small.

- **Sex:** From 2010 through 2014, the rates for males and for females overall decreased (Table 1a). In 2014, males accounted for 74% of all diagnosed HIV infections. The rate for males (10.7) was 3 times the rate for females (3.4) (Table 1a). Among males, the rates for those aged 50–64 years decreased, but rates for those aged 65 years and older remained stable. Among females, the rates for those aged 50–64 years decreased, but the rates for females aged 65 years and older remained stable (Table 2a).
- **Transmission category:** From 2010 through 2014, among males, the estimated annual number of diagnosed HIV infections attributed to male-to-male sexual contact increased. The estimated numbers of infections attributed to heterosexual contact, injection drug use, or male-to-male sexual contact *and* injection drug use decreased. Among females, the estimated number of diagnosed infections attributed to heterosexual contact remained stable, but the number of infections attributed to injection drug use decreased. In 2014, among males, 67% of diagnosed infections were attributed to male-to-male sexual contact, 20% to heterosexual contact, 10% to injection drug use, and 3% to male-to-male sexual contact *and* injection drug use; among females, 85% of diagnosed infections were attributed to heterosexual contact and 15% to injection drug use (Table 1a).

Adults Aged 50 Years and Older Living with Diagnosed HIV Infection (Prevalence)

From 2010 through 2013, the estimated number of older adults living with diagnosed HIV infection (prevalence) increased from 301,970 (rate, 303.1) at the end of 2010 to 395,668 (rate, 371.0) at the end of 2013 (Table 6a).

- **Age group:** At the end of 2013, the largest percentage of older adults living with diagnosed HIV infection was among those aged 50–54 years (41%), followed by those aged 55–59 years (29%). From 2010 through 2013, prevalence rates increased among older adults in all age groups (Table 6a).

- **Race/ethnicity:** At the end of 2013, the largest percentage of older adults living with diagnosed HIV (40%) was that among blacks/African Americans. Among the remaining race/ethnicity groups, the percentages were 38%, whites; 17%, Hispanics/Latinos; 3%, persons of multiple races; 1%, Asians; and less than 1% each, American Indians/Alaska Natives and Native Hawaiians/ other Pacific Islanders. Among all race/ethnicity groups, the highest estimated prevalence rate of diagnosed HIV infection was that for blacks/African Americans: 1,453.8 per 100,000 population. The prevalence rate among blacks/African Americans was 2 times the rate for Hispanics/Latinos (685.9) and almost 8 times the rate for whites (190.1) (Table 6a).
- **Sex and transmission category:** At the end of 2013, among an estimated total of 306,722 males living with diagnosed HIV infection, 62% of infections were attributed to male-to-male sexual contact, 17% to injection drug use, 12% to heterosexual contact, and 8% to male-to-male sexual contact *and* injection drug use. Among an estimated total of 88,946 females living with diagnosed HIV infection, 69% of infections were attributed to heterosexual contact and 30% to injection drug use. The estimated prevalence rates per 100,000 population were 617.7 for males and 156.0 for females (Table 6a).
- **Population of area of residence:** At the end of 2013, the prevalence rate of older adults living with diagnosed HIV infection in metropolitan statistical areas (482.4) was 3 times the rate in metropolitan areas (189.9) and 4 times the rate in nonmetropolitan areas (116.2). The prevalence rates were highest for adults of multiple races (1,759.9) and blacks/African Americans (1,628.8) residing in metropolitan statistical areas (Table 11).

Deaths of Adults Aged 50 Years and Older with Diagnosed HIV Infection

From 2010 through 2013, the annual estimated number of deaths of older adults with diagnosed HIV infection increased; the annual rate of deaths remained stable. In 2013, the estimated rate of deaths of older adults with diagnosed HIV infection was 9.5 per 100,000 population. Deaths of persons

with diagnosed HIV infection may be due to any cause (i.e., may or may not be related to HIV infection) (Table 12a).

- **Age group:** From 2010 through 2013, the rates of deaths for adults aged 65–69, 70–74, 75–79, and 80–84 years increased. The rate for adults aged 50–54 years decreased, and rates of deaths in all other age groups remained stable (Table 12a). In 2013, the rate of deaths for adults aged 50–64 years (12.7) with diagnosed HIV infection was 2 times the rate for adults aged 65 years and older (5.1) (Table 13a).
- **Race/ethnicity:** From 2010 through 2013, the rates of deaths for whites and American Indians/Alaska Natives increased. The rates of deaths for Asians, blacks/African Americans, and persons of multiple races decreased. The rates of deaths for Hispanics/Latinos remained stable. In 2013, the highest rate of deaths was for persons of multiple races (57.0). The rate of deaths among blacks/African Americans (42.6) was 3 times the rate among Hispanics/Latinos (15.6) and 10 times the rate among whites (4.2) (Table 12a). Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** From 2010 through 2013, the rates of deaths for males remained stable. The rates of deaths for females increased. The estimated number of deaths among males with infection attributed to male-to-male sexual contact or male-to-male sexual contact *and* injection drug use increased. The estimated number of deaths among females with infection attributed to injection drug use or heterosexual contact increased. The rate of deaths among males (15.6) was 4 times the rate among females (4.2) (Table 12a).

Stage 3 (AIDS) Classification at the Time of HIV Diagnosis

Among older adults with HIV infection diagnosed during 2013, 37.7% of infections were classified as stage 3 (AIDS) at the time of diagnosis (Table 15a).

- **Age group:** The percentage of older adults with stage 3 (AIDS) classification at the time of diagnosis was 36.8% among adults aged 50–54 years and was higher among older age groups, with the

exception of adults aged 80–84 years and 85 years and older (Table 15a). Please use caution when interpreting data for adults aged 85 years and older: the numbers are small.

- **Race/ethnicity:** The highest percentage of older adults with stage 3 (AIDS) classification at the time of diagnosis was among Asians (51.6%), followed by the percentages among persons of multiple races (46.0%), Hispanics/Latinos (41.6%), whites (39.2%), American Indians/Alaska Natives (37.0%), and blacks/African Americans (34.2%) (Table 15a). Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** Among males, the percentage with stage 3 (AIDS) classification at the time of diagnosis was highest among those with infection attributed to heterosexual contact (41.4%). Among females, the percentage was highest among those with infection attributed to injection drug use (36.8%) (Table 15a).

SUGGESTED READINGS

- Brooks JT, Buchacz K, Gebo KA, Mermin J. HIV infection and older Americans: the public health perspective. *Am J Public Health* 2012;102(8):1516–1526. doi:10.2105/AJPH.2012.300844.
- CDC. Establishing a holistic framework to reduce inequities in HIV, viral hepatitis, STDs, and tuberculosis in the United States: an NCHHSTP white paper on social determinants of health, 2010. <http://www.cdc.gov/socialdeterminants/docs/SDH-White-Paper-2010.pdf>. Published October 2010. Accessed April 25, 2016.
- CDC. Estimated HIV incidence in the United States, 2007–2010. *HIV Surveillance Supplemental Report* 2012;17(No. 4). <http://go.usa.gov/p8P4>. Published December 2012. Accessed April 25, 2016.
- CDC. *HIV Surveillance Report, 2014*; vol. 26. <http://www.cdc.gov/hiv/library/reports/surveillance/>. Published November 2015. Accessed April 25, 2016.
- CDC. HIV surveillance—United States, 1981–2008. *MMWR* 2011;60(21):689–693.

- CDC. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 dependent areas, 2013. *HIV Surveillance Supplemental Report* 2015;20(No. 2). <http://www.cdc.gov/hiv/library/reports/surveillance/>. Published July 2015. Accessed April 25, 2016.
- CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.
- CDC. Revised surveillance case definitions for HIV infection—United States, 2014. *MMWR* 2014;63(RR-03):1–10.
- CDC. Terms, definitions, and calculations used in CDC HIV surveillance publications. <http://www.cdc.gov/hiv/statistics/recommendations/terms.html>. Published September 2014. Accessed April 25, 2016.
- CDC. Vital Signs: HIV diagnosis, care, and treatment among persons living with HIV—United States, 2011. *MMWR* 2014;63(47):1113–1117.
- CDC. Vital Signs: HIV prevention through care and treatment—United States. *MMWR* 2011;60(47):1618–1623.
- CDC. Vital Signs: HIV testing and diagnosis among adults—United States. *MMWR* 2010;59(47):1550–1555.
- Cohen SM, Gray KM, Ocfemia MC, Johnson AS, Hall HI. The status of the National HIV Surveillance System, United States, 2013. *Public Health Rep* 2014;129(4):335–341.
- Hall HI, Song R, Rhodes P, et al. Estimation of HIV incidence in the United States. *JAMA* 2008;300(5):520–529. doi: 10.1001/jama.300.5.520.
- Johnson AS, Hall HI, Hu X, Lansky A, Holtgrave DR, Mermin J. Trends in diagnoses of HIV infection in the United States, 2002–2011 [Research Letter]. *JAMA* 2014;312(4):432–434. doi:10.1001/jama.2014.8534.
- Linley L, Prejean J, An Q, Chen M, Hall HI. Racial/ethnic disparities in HIV diagnoses among persons aged 50 years and older in 37 US states, 2005–2008. *Am J Public Health* 2012;102(8):1527–1534. doi:10.2105/AJPH.2011.300431.
- Prejean J, Song R, Hernandez A, et al. Estimated HIV incidence in the United States, 2006–2009. *PLoS One* 2011;6(8):e17502. doi:10.1371/journal.pone.0017502.

SURVEILLANCE OF HIV INFECTION

This report includes data from case reports from 50 states, the District of Columbia, and 6 dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting to the jurisdiction (not to CDC), by name, for persons with confirmed diagnoses of HIV infection. After the removal of personally identifiable information, data from these reports were submitted to CDC.

TABULATION AND PRESENTATION OF DATA

The data in this report include information received by CDC through July 31, 2015. Data are presented for adults aged 50 years and older whose HIV infection was diagnosed during 2010–2014. Data are also presented for persons living with diagnosed HIV infection and for deaths among persons with diagnosed HIV infection as of December 31, 2013. Data on stage 3 (AIDS) classifications at the time of initial HIV diagnosis are also presented for persons whose HIV infection was diagnosed during 2013.

Data tables are organized as follows.

- Diagnosed HIV infection among adults aged 50 years and older, by selected characteristics (Tables 1a/b–5)
- Adults aged 50 years and older living with diagnosed HIV infection, by selected characteristics (Tables 6a/b–11)
- Deaths among adults aged 50 years and older with diagnosed HIV infection, by selected characteristics (Tables 12 a/b–14a/b)
- Stage 3 (AIDS) classification at the time of HIV diagnosis among adults aged 50 years and older, by selected characteristics (Tables 15 a/b–16a/b)

Please use caution when interpreting reported numbers less than 12, as well as estimated numbers and accompanying rates based on these numbers: the underlying relative standard errors are more than 30%, so the numbers are considered unreliable.

Diagnoses of HIV Infection

Data in this report were compiled from diagnoses of HIV infection that were reported by the United States

and 6 dependent areas to CDC through July 31, 2015. All jurisdictions report diagnoses of HIV infection to CDC on the basis of guidelines for HIV case surveillance. Both the 2008 and 2014 HIV case definitions were used to classify HIV infection (see Stages of HIV Infection below) [2, 3]. The 2008 case definition was used to classify infections diagnosed from 2010 through 2013. The 2014 case definition was used to classify infections diagnosed in 2014. Please use caution when interpreting data on diagnoses of HIV infection. HIV surveillance reports may not be representative of all persons with HIV because not all infected persons have (1) been tested or (2) tested at a time when the infection could be detected and diagnosed. Also, some states offer anonymous HIV testing; the results of anonymous tests are not reported to the confidential name-based HIV registries of state and local health departments. Therefore, reports of confidential test results may not represent all persons who tested positive for HIV infection. In addition, testing patterns are influenced by many factors, including the extent to which testing is routinely offered to specific groups, or the availability of, and access to, medical care and testing services. The data presented in this report provide a minimum estimate of persons for whom HIV infection has been diagnosed and reported to the surveillance system.

Persons Living with Diagnosed HIV Infection (Prevalence)

Prevalence of HIV infection was based on the number of persons living with diagnosed HIV infection, regardless of the stage of disease, as of December 31, 2013. The exclusion of data from the most recent year allowed at least 18 months for deaths to be reported and for these deaths to be factored into calculations of prevalence.

Deaths

Persons reported to the National HIV Surveillance System are assumed alive unless their deaths have been reported to CDC. Death data include all deaths of persons with diagnosed HIV infection regardless of the cause of death (Tables 12a/b–14a/b). Data are reported for deaths that occurred by December 31, 2013, thereby allowing at least 18 months for deaths

to be reported to CDC. The estimated numbers and rates of deaths resulted from statistical adjustment for delays in reporting.

Stages of HIV Infection

Both the 2008 and 2014 HIV case definitions were used to classify HIV infection among older adults [2, 3]. The 2008 case definition was used to classify cases diagnosed from 2010 through 2013. This definition incorporates an HIV infection staging system that includes AIDS (HIV infection, stage 3). The 2008 stages of HIV infection are defined as follows:

- **HIV infection, stage 1:** No AIDS-defining condition and either CD4 count of ≥ 500 cells/ μL or CD4 percentage of total lymphocytes of ≥ 29 .
- **HIV infection, stage 2:** No AIDS-defining condition and either CD4 count of 200–499 cells/ μL or CD4 percentage of total lymphocytes of 14–28.
- **HIV infection, stage 3 (AIDS):** Documentation of an AIDS-defining condition or either a CD4 count of < 200 cells/ μL or CD4 percentage of total lymphocytes of < 14 . Documentation of an AIDS-defining condition supersedes a CD4 count or percentage that would not, by itself, be the basis for a stage 3 (AIDS) classification.
- **HIV infection, stage unknown:** No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

The 2014 case definition was used to classify cases diagnosed in 2014. It is similar to the 2008 case definition except for the following: (1) inclusion of criteria for stage 0, (2) the inclusion of CD4 testing criteria for stage 3 in children, and (3) changes in the cutoffs for CD4 percentage of total lymphocytes used for classification of stages 1 and 2 in persons aged 6 years and older. The stages of HIV infection in the 2014 case definition are based on age-specific CD4+ T-lymphocyte counts or percentages of total lymphocytes and are defined as follows:

- **HIV infection, stage 0:** First positive HIV test result within 6 months after a negative HIV test result. The stage remains stage 0 until 6 months after the first positive test result. After 6 months, the stage may be reclassified as 1, 2, 3, or unknown if based on a CD4+ T-lymphocyte result or the diagnosis of an OI [opportunistic illness]. The diagnosis of an AIDS-defining condition or a low CD4 test result before the 6 months

have elapsed does not change the stage from stage 0 to stage 3.

- **HIV infection, stages 1, 2, and 3:** Documentation of an AIDS-defining condition (excluding stage 0 as described above) is stage 3. Otherwise, the stage is determined by the lowest CD4 test result [3, p 7].
- **HIV infection, stage unknown:** No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

In this report, the term *diagnosis of HIV infection* is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 0, 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. The term *HIV infection, stage 3 (AIDS)*, and its condensed version—*stage 3 (AIDS)*—refer specifically to persons with diagnosed HIV whose infection was classified as stage 3 (AIDS) during a given year (for diagnoses) or whose infection has ever been classified as stage 3 (AIDS) (for prevalence and deaths).

Age

For presentations of data on older adults living with diagnosed HIV infection (Tables 6a/b–11), the age-group assignment (for example, 50–64 years) is based on the person's age as of December 31 of the specified year. In Tables 12a/b–14a/b, which present data on deaths of persons with a diagnosis of HIV infection or whose infection has ever been classified as stage 3 (AIDS), age-group assignment is determined by the person's age at the time of death. For all other tables, the age-group assignment is based on the person's age at the time of diagnosis of HIV infection.

Sex

Sex designations in this report are based on a person's sex at birth. *Transgender* is an umbrella term that is used to identify persons whose sex assigned at birth does not match current gender identity or expression. Data for transgender persons are not presented in this report because information on gender identity (a person's internal understanding of his or her gender or the gender with which a person identifies) is not consistently collected or documented in the data sources used by HIV reporting jurisdictions. HIV surveillance personnel collect this information, when available, from sources such as case report forms submitted by health care or HIV testing providers and medical

records, or by matching with other databases (e.g., Ryan White program data) in the health department. In May 2012, CDC issued guidance to state and local programs on methods for collecting data on transgender persons and working with transgender-specific data. However, characterization of HIV infection among transgender persons may require supplemental data from special studies.

Race and Ethnicity

In the *Federal Register* for October 30, 1997 [4], the Office of Management and Budget (OMB) announced the Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity. Implementation by January 1, 2003, was mandated. At a minimum, data on the following race categories should be collected:

- American Indian or Alaska Native
- Asian
- black or African American
- Native Hawaiian or other Pacific Islander
- white

Additionally, systems must be able to retain information when multiple race categories are reported. In addition to data on race, data on 2 categories of ethnicity should be collected:

- Hispanic or Latino
- not Hispanic or Latino

The Asian or Pacific Islander category in annual surveillance reports published prior to the 2007 surveillance report was split into 2 categories: (1) Asian and (2) Native Hawaiian or other Pacific Islander. The Asian category (in tables where footnoted) includes the cases in Asians/Pacific Islanders (referred to as legacy cases) that were reported before the implementation of the new race categories in 2003 (e.g., cases of HIV infection that were diagnosed and reported to CDC before 2003 but that progressed to AIDS after 2003) and a small percentage of cases that were reported after 2003 but that were reported according to the old race category (Asian/Pacific Islander). In tables of diagnoses of HIV infection during 2010–2014, the Asian category does not include Asian/Pacific Islander cases because these cases were diagnosed after 2003 and were reported to CDC in accordance with OMB's Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity [4].

This report also presents data for persons for whom multiple race categories are reported. In this report, persons categorized by race were not Hispanic or Latino. The number of persons reported in each race category may, however, include persons whose ethnicity was not reported.

Population of Area of Residence

In the *Federal Register* for June 28, 2010, OMB published revised standards for defining metropolitan statistical areas in federal statistical activities [5]. These standards, which provided for the identification of metropolitan statistical areas in the United States and Puerto Rico, replaced the 2000 standards. The adoption of the new standards was effective as of June 28, 2010. On February 28, 2013, OMB announced new metropolitan statistical area definitions based on the new standards and Census 2010 data [6]. In this report, populations of area of residence are defined as areas with populations of 500,000 or more (metropolitan statistical areas), areas with populations of 50,000–499,999 (metropolitan areas), or areas with populations less than 50,000 (nonmetropolitan areas).

Transmission Categories

Transmission category is the term for the classification of cases that summarizes a person's possible HIV risk factors; the summary classification results from selecting, from the presumed hierarchical order of probability, the 1 (single) risk factor most likely to have been responsible for transmission. For surveillance purposes, a diagnosis of HIV infection is counted only once in the hierarchy of transmission categories. Persons with more than 1 reported risk factor for HIV infection are classified in the transmission category listed first in the hierarchy. The exception is men who had sexual contact with other men *and* injection drug use; this group makes up a separate transmission category.

Persons whose transmission category is classified as male-to-male sexual contact include men who have ever had sexual contact with other men (i.e., homosexual contact) and men who have ever had sexual contact with both men and women (i.e., bisexual contact). Persons whose transmission category is classified as heterosexual contact are persons who report heterosexual contact with a person known to have, or to be at high risk for, HIV infection (e.g., an injection drug user).

Cases of HIV infection reported without a risk factor listed in the hierarchy of transmission categories are classified as “no identified risk (NIR)” [7]. Cases classified as NIR include cases that are being followed up by local health department staff; cases in persons whose risk-factor information is missing because they died, declined to be interviewed, or were lost to follow-up; and cases in persons who were interviewed or for whom other follow-up information was available but for whom no risk factor was identified.

Because a substantial proportion of cases of HIV infection are reported to CDC without an identified risk factor, multiple imputation is used to assign a transmission category [8]. Multiple imputation is a statistical approach in which each missing transmission category is replaced with a set of plausible values that represent the uncertainty about the true, but missing, value [9]. The plausible values are analyzed by using standard procedures, and the results from these analyses are then combined to produce the final results.

Estimation Methods

Diagnoses of HIV infection in 2010 through 2014 and prevalence of HIV infection at year-end 2013 were estimated by using the same methods that were used for the 2014 *HIV Surveillance Report* [1]. Estimated numbers and rates were not adjusted for incomplete reporting. Estimated numbers were rounded to zero decimal places and rates to one decimal place.

Reporting Delays

Reporting delays (time between diagnosis or death and the reporting of diagnosis or death to CDC) may differ among demographic and geographic categories; for some, delays in reporting have been as long as several years. The statistical adjustment of the data on diagnoses and deaths is based on estimates of reporting-delay distributions, which are calculated by using a modified semiparametric life-table statistical procedure. This procedure takes into account differences in reporting delays due to sex, race/ethnicity, HIV transmission categories, geographic area (i.e., reporting city, state, or dependent area; region of residence), the size of the metropolitan statistical area, and the type of facility where the diagnosis was made or death occurred [10]. The statistical adjustment of data on diagnoses takes into consideration the infor-

mation reported from the state or dependent area where the diagnosis was made; the statistical adjustment of data on deaths takes into consideration the information from the state where the death occurred.

Rates

Rates per 100,000 population were calculated for (1) the numbers of diagnoses of HIV infection, (2) the numbers of persons living with diagnosed HIV infection, and (3) the numbers of deaths of persons with diagnosed HIV infection.

The population denominators used to compute the rates for the 50 states, the District of Columbia, and Puerto Rico were based on the Vintage 2014 postcensal estimates file (for years 2010–2014) from the U.S. Census Bureau [11]. The population denominators for American Samoa, Guam, the Northern Mariana Islands, the Republic of Palau, and the U.S. Virgin Islands were based on estimates and projections from the U.S. Census Bureau’s International Data Base [12]. Each rate was calculated by dividing the estimated total number of diagnoses (or deaths, or persons living with diagnosed HIV infection) for the calendar year by the population for that calendar year and then multiplying the number by 100,000. The denominators used for calculating the rates specific to age, sex, and race/ethnicity were computed by applying the appropriate vintage estimates for age, sex, and race/ethnicity for the 50 states and the District of Columbia [11]. The same method was used to calculate the denominators for Puerto Rico, with the exception of race/ethnicity estimates; these data are not available for Puerto Rico (see next paragraph). For the other 5 U.S. dependent areas, estimates from the U.S. Census Bureau’s International Data Base were used for age- and sex-specific population denominators [12].

CDC currently does not provide subpopulation rates by race/ethnicity for the 6 dependent areas because the U.S. Census Bureau does not collect information from all dependent areas. Rates for transmission categories are not provided because of the absence of denominator data from the U.S. Census Bureau, the source of data used for calculating all rates in this report.

References

1. CDC. *HIV Surveillance Report, 2014*; vol. 26. <http://www.cdc.gov/hiv/library/reports/surveillance/>. Published November 2015. Accessed April 25, 2016.
2. CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.
3. CDC. Revised surveillance case definition for HIV infection—United States, 2014. *MMWR* 2014;63(RR-03):1–10.
4. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Federal Register* 1997;62:58781–58790. <http://go.usa.gov/vSdR>. Accessed April 25, 2016.
5. Office of Management and Budget. 2010 Standards for delineating metropolitan and micropolitan statistical areas. *Federal Register* 2010;75(123):37246–39052. <http://go.usa.gov/vSpG>. Accessed April 25, 2016.
6. Office of Management and Budget. Revised delineations of metropolitan statistical areas, micropolitan statistical areas, and combined statistical areas, and guidance on uses of the delineations of these areas. OMB Bulletin 13-01. <http://go.usa.gov/vSyB>. Published February 28, 2013. Accessed April 25, 2016.
7. CDC. Current Trends: Heterosexually acquired AIDS—United States, 1993. *MMWR* 1994;43(9):158, Fig 3.
8. McDavid Harrison K, Kajese T, Hall HI, Song R. Risk factor redistribution of the national HIV/AIDS surveillance data: an alternative approach. *Public Health Rep* 2008;123(5):618–627.
9. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. New York: John Wiley & Sons Inc; 1987.
10. Song R, Green TA. An improved approach to accounting for reporting delay in case surveillance systems. *JP J Biostat* 2012;7(1):1–14.
11. U.S. Census Bureau. Population estimates [entire data set]. <http://www.census.gov/popest/data/>. Updated July 1, 2014. Accessed April 25, 2016.
12. U.S. Census Bureau. International Data Base. <http://go.usa.gov/vSUj>. Updated December 2013. Accessed April 25, 2016.

Table 1a. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States

	2010			2011			2012			2013			2014		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate												
Males															
Age at diagnosis (yr)															
50–54	2,543	2,608	23.8	2,487	2,578	23.3	2,414	2,532	22.8	2,208	2,359	21.3	2,228	2,464	22.2
55–59	1,472	1,511	15.8	1,461	1,514	15.4	1,353	1,421	14.1	1,455	1,553	15.1	1,369	1,521	14.6
60–64	773	792	9.7	757	784	9.2	745	782	9.2	795	847	9.8	690	766	8.6
65–69	329	338	5.7	369	383	6.3	380	399	6.0	393	419	6.1	377	420	5.8
70–74	153	157	3.7	159	165	3.7	134	140	3.1	156	167	3.4	135	150	2.9
75–79	67	69	2.2	66	68	2.1	61	64	2.0	59	64	1.9	73	81	2.3
80–84	19	20	0.8	18	19	0.8	18	19	0.8	26	28	1.2	23	26	1.1
≥85	11	11	0.6	3	3	0.2	11	12	0.6	11	12	0.6	3	3	0.1
Race/ethnicity															
American Indian/Alaska Native	20	20	7.6	16	16	5.9	14	15	5.1	15	16	5.2	17	19	6.0
Asian	64	66	3.7	66	69	3.7	68	71	3.6	56	60	2.9	79	88	4.1
Black/African American	2,206	2,266	52.1	2,107	2,191	48.7	1,970	2,078	44.7	1,815	1,951	40.7	1,732	1,938	39.3
Hispanic/Latino ^b	789	808	20.3	830	858	20.4	817	856	19.2	842	898	19.1	812	896	18.1
Native Hawaiian/Other Pacific Islander	2	2	4.1	3	3	6.1	7	7	13.2	2	2	3.5	4	4	7.2
White	2,105	2,156	6.1	2,151	2,225	6.2	2,125	2,223	6.1	2,255	2,398	6.4	2,171	2,393	6.3
Multiple races	181	187	50.8	147	152	39.6	115	121	30.0	118	125	29.7	83	94	21.5
Transmission category															
Male-to-male sexual contact	2,236	3,257	—	2,411	3,431	—	2,310	3,415	—	2,356	3,528	—	2,338	3,654	-
Injection drug use	460	759	—	376	643	—	322	593	—	291	554	—	257	537	-
Male-to-male sexual contact and injection drug use	140	194	—	135	186	—	113	165	—	108	163	—	101	155	-
Heterosexual contact ^c	763	1,275	—	764	1,239	—	702	1,174	—	673	1,188	—	572	1,069	-
Other ^d	1,768	21	—	1,634	15	—	1,669	24	—	1,675	17	—	1,630	17	-
Subtotal, males	5,367	5,505	11.9	5,320	5,514	11.6	5,116	5,370	11.1	5,103	5,450	11.0	4,898	5,432	10.7
Females															
Age at diagnosis (yr)															
50–54	967	993	8.7	877	909	7.9	805	845	7.3	816	876	7.6	700	778	6.8
55–59	605	621	6.1	538	558	5.3	570	598	5.6	592	631	5.8	580	645	5.8
60–64	291	299	3.4	315	327	3.5	307	324	3.5	303	324	3.4	270	303	3.1
65–69	146	150	2.3	116	121	1.8	120	125	1.7	121	130	1.7	136	153	1.9
70–74	34	35	0.7	53	55	1.1	55	59	1.1	64	70	1.2	49	55	0.9
75–79	18	18	0.4	22	23	0.6	24	25	0.6	27	29	0.7	15	17	0.4
80–84	11	11	0.3	7	7	0.2	11	12	0.3	8	8	0.2	3	4	0.1
≥85	2	2	0.1	3	3	0.1	5	5	0.1	2	2	0.1	5	6	0.1

Table 1a. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States (cont)

	2010			2011			2012			2013			2014		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate												
Race/ethnicity															
American Indian/Alaska Native	12	12	4.0	8	8	2.6	6	6	1.9	15	16	4.7	10	11	3.0
Asian	14	14	0.7	24	25	1.1	16	17	0.7	19	20	0.8	23	26	1.0
Black/African American	1,349	1,386	24.8	1,269	1,319	22.8	1,249	1,315	22.1	1,215	1,304	21.3	1,118	1,250	19.9
Hispanic/Latino ^b	302	310	6.8	251	260	5.4	257	270	5.3	292	311	5.8	278	309	5.5
Native Hawaiian/Other Pacific Islander	0	0	0.0	2	2	3.8	1	1	1.7	4	4	6.6	2	2	3.2
White	328	335	0.8	321	332	0.8	313	326	0.8	337	359	0.9	305	337	0.8
Multiple races	69	71	16.7	56	58	13.1	55	58	12.4	51	55	11.4	22	24	4.7
Transmission category															
Injection drug use	215	399	—	174	344	—	170	335	—	161	320	—	121	288	-
Heterosexual contact ^c	933	1,723	—	893	1,653	—	836	1,650	—	898	1,743	—	731	1,667	-
Other ^d	926	7	—	864	7	—	891	8	—	874	7	—	906	4	-
Subtotal, females	2,074	2,130	4.0	1,931	2,004	3.7	1,897	1,993	3.6	1,933	2,070	3.6	1,758	1,959	3.4
Both sexes															
Age at diagnosis (yr)															
50–54	3,510	3,602	16.1	3,364	3,487	15.4	3,219	3,377	14.9	3,024	3,235	14.3	2,928	3,242	14.4
55–59	2,077	2,132	10.8	1,999	2,072	10.2	1,923	2,019	9.7	2,047	2,184	10.3	1,949	2,166	10.1
60–64	1,064	1,091	6.4	1,072	1,111	6.2	1,052	1,106	6.2	1,098	1,170	6.5	960	1,069	5.8
65–69	475	487	3.9	485	504	3.9	500	524	3.7	514	549	3.8	513	573	3.7
70–74	187	191	2.0	212	220	2.3	189	199	2.0	220	237	2.2	184	205	1.9
75–79	85	87	1.2	88	92	1.2	85	90	1.2	86	93	1.2	88	98	1.2
80–84	30	31	0.5	25	26	0.4	29	31	0.5	34	37	0.6	26	30	0.5
≥85	13	13	0.2	6	6	0.1	16	17	0.3	13	14	0.2	8	9	0.1
Race/ethnicity															
American Indian/Alaska Native	32	33	5.7	24	25	4.1	20	21	3.4	30	32	4.9	27	29	4.4
Asian	78	80	2.0	90	93	2.2	84	88	2.0	75	80	1.7	102	114	2.3
Black/African American	3,555	3,652	36.7	3,376	3,509	34.1	3,219	3,393	32.0	3,030	3,255	29.8	2,850	3,189	28.4
Hispanic/Latino ^b	1,091	1,118	13.1	1,081	1,118	12.4	1,074	1,125	11.8	1,134	1,209	12.1	1,090	1,205	11.4
Native Hawaiian/Other Pacific Islander	2	2	2.0	5	5	4.9	8	8	7.2	6	6	5.1	6	7	5.1
White	2,433	2,492	3.3	2,472	2,557	3.3	2,438	2,549	3.3	2,592	2,758	3.5	2,476	2,730	3.4
Multiple races	250	258	32.5	203	211	25.4	170	178	20.6	169	180	19.9	105	118	12.5
Total^e	7,441	7,635	7.7	7,251	7,518	7.4	7,013	7,363	7.1	7,036	7,520	7.1	6,656	7,391	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 1b. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States and 6 dependent areas

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
Males															
Age at diagnosis (yr)															
50–54	2,597	2,665	24.0	2,553	2,647	23.6	2,461	2,582	23.0	2,262	2,418	21.6	2,262	2,504	22.4
55–59	1,504	1,544	15.9	1,496	1,551	15.6	1,377	1,447	14.2	1,481	1,581	15.2	1,400	1,558	14.8
60–64	791	811	9.8	776	804	9.3	758	796	9.2	822	876	10.0	712	792	8.8
65–69	342	351	5.9	376	391	6.3	388	407	6.1	405	432	6.2	383	427	5.8
70–74	157	161	3.7	162	168	3.8	139	146	3.1	159	171	3.4	139	155	3.0
75–79	69	71	2.2	69	71	2.2	69	73	2.2	62	67	2.0	75	84	2.3
80–84	20	21	0.9	18	19	0.8	20	21	0.9	29	32	1.3	23	26	1.1
≥85	13	14	0.7	3	3	0.2	12	13	0.6	11	12	0.6	3	3	0.1
Race/ethnicity															
American Indian/Alaska Native	20	20	—	16	16	—	14	15	—	15	16	—	17	19	—
Asian	64	66	—	66	69	—	68	71	—	56	60	—	80	89	—
Black/African American	2,209	2,269	—	2,112	2,196	—	1,970	2,078	—	1,818	1,954	—	1,738	1,945	—
Hispanic/Latino ^b	912	936	—	958	993	—	925	972	—	967	1,035	—	902	1,001	—
Native Hawaiian/Other Pacific Islander	2	2	—	3	3	—	7	7	—	2	2	—	4	4	—
White	2,105	2,156	—	2,151	2,225	—	2,125	2,223	—	2,255	2,398	—	2,172	2,394	—
Multiple races	181	187	—	147	152	—	115	121	—	118	125	—	84	95	—
Transmission category															
Male-to-male sexual contact	2,264	3,288	—	2,449	3,474	—	2,340	3,447	—	2,385	3,562	—	2,368	3,693	—
Injection drug use	488	794	—	400	675	—	343	618	—	319	589	—	273	563	—
Male-to-male sexual contact and injection drug use	143	198	—	138	189	—	116	168	—	113	168	—	102	157	—
Heterosexual contact ^c	816	1,335	—	818	1,300	—	752	1,229	—	729	1,253	—	609	1,117	—
Other ^d	1,782	21	—	1,648	15	—	1,673	24	—	1,685	17	—	1,645	17	—
Subtotal, males	5,493	5,636	12.0	5,453	5,654	11.8	5,224	5,486	11.2	5,231	5,590	11.1	4,997	5,547	10.8
Females															
Age at diagnosis (yr)															
50–54	983	1,010	8.8	899	932	8.0	831	872	7.5	842	904	7.8	711	791	6.8
55–59	619	636	6.1	551	572	5.4	583	612	5.6	611	652	5.9	591	658	5.9
60–64	309	318	3.5	321	334	3.6	317	335	3.6	312	334	3.5	274	307	3.1
65–69	152	156	2.3	123	128	1.9	132	138	1.8	125	135	1.7	137	154	1.9
70–74	35	36	0.7	55	57	1.1	56	60	1.1	64	70	1.2	52	58	1.0
75–79	18	18	0.4	23	25	0.6	25	27	0.6	27	29	0.7	15	17	0.4
80–84	11	11	0.3	9	9	0.3	11	12	0.3	9	10	0.3	3	4	0.1
≥85	2	2	0.1	3	3	0.1	5	5	0.1	2	2	0.1	5	6	0.1

Table 1b. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, sex, and selected characteristics, 2010–2014—United States and 6 dependent areas (cont)

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
Race/ethnicity															
American Indian/Alaska Native	12	12	—	8	8	—	6	6	—	15	16	—	10	11	—
Asian	14	14	—	24	25	—	16	17	—	19	20	—	23	26	—
Black/African American	1,350	1,387	—	1,269	1,319	—	1,249	1,315	—	1,217	1,307	—	1,120	1,252	—
Hispanic/Latino ^b	356	367	—	304	316	—	320	337	—	348	373	—	306	341	—
Native Hawaiian/Other Pacific Islander	0	0	—	2	2	—	1	1	—	4	4	—	2	2	—
White	328	335	—	321	332	—	313	326	—	338	360	—	305	337	—
Multiple races	69	71	—	56	58	—	55	58	—	51	55	—	22	24	—
Transmission category															
Injection drug use	217	401	—	178	349	—	175	342	—	170	331	—	123	291	—
Heterosexual contact ^c	983	1,778	—	938	1,704	—	890	1,711	—	937	1,797	—	757	1,699	—
Other ^d	929	7	—	868	7	—	895	8	—	885	7	—	908	4	—
Subtotal, females	2,129	2,187	4.0	1,984	2,060	3.7	1,960	2,061	3.6	1,992	2,135	3.7	1,788	1,994	3.4
Both sexes															
Age at diagnosis (yr)															
50–54	3,580	3,674	16.2	3,452	3,580	15.7	3,292	3,455	15.1	3,104	3,323	14.5	2,973	3,294	14.4
55–59	2,123	2,179	10.9	2,047	2,123	10.3	1,960	2,059	9.8	2,092	2,234	10.4	1,991	2,215	10.2
60–64	1,100	1,129	6.6	1,097	1,138	6.3	1,075	1,131	6.3	1,134	1,210	6.6	986	1,099	5.8
65–69	494	507	4.0	499	519	4.0	520	546	3.8	530	567	3.8	520	581	3.7
70–74	192	196	2.1	217	226	2.3	195	206	2.0	223	240	2.2	191	213	1.9
75–79	87	90	1.2	92	96	1.3	94	99	1.3	89	96	1.2	90	100	1.2
80–84	31	32	0.5	27	28	0.5	31	33	0.6	38	41	0.7	26	30	0.5
≥85	15	16	0.3	6	6	0.1	17	18	0.3	13	14	0.2	8	9	0.1
Race/ethnicity															
American Indian/Alaska Native	32	33	—	24	25	—	20	21	—	30	32	—	27	29	—
Asian	78	80	—	90	93	—	84	88	—	75	80	—	103	115	—
Black/African American	3,559	3,656	—	3,381	3,515	—	3,219	3,393	—	3,035	3,261	—	2,858	3,198	—
Hispanic/Latino ^b	1,268	1,303	—	1,262	1,309	—	1,245	1,309	—	1,315	1,407	—	1,208	1,342	—
Native Hawaiian/Other Pacific Islander	2	2	—	5	5	—	8	8	—	6	6	—	6	7	—
White	2,433	2,492	—	2,472	2,557	—	2,438	2,549	—	2,593	2,759	—	2,477	2,731	—
Multiple races	250	258	—	203	211	—	170	178	—	169	180	—	106	119	—
Total^e	7,622	7,823	7.8	7,437	7,714	7.5	7,184	7,546	7.1	7,223	7,725	7.2	6,785	7,541	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2a. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
50–64 years of age at diagnosis															
Males															
Race/ethnicity															
American Indian/Alaska Native	16	16	8.7	14	14	7.4	13	14	6.9	12	13	6.2	16	18	8.6
Asian	58	60	5.1	59	61	5.1	59	62	4.9	50	54	4.1	65	72	5.3
Black/African American	1,989	2,044	67.9	1,864	1,938	62.2	1,729	1,824	57.2	1,577	1,695	52.0	1,490	1,668	50.2
Hispanic/Latino ^b	702	719	25.8	728	752	25.5	726	760	24.5	757	807	24.8	729	805	23.5
Native Hawaiian/Other Pacific Islander	2	2	5.7	3	3	8.6	6	6	16.2	2	2	5.1	3	3	8.1
White	1,858	1,903	8.9	1,906	1,971	9.1	1,874	1,959	9.0	1,958	2,082	9.6	1,909	2,102	9.7
Multiple races	163	168	65.7	131	136	50.8	105	110	40.0	102	107	37.7	75	85	28.9
Transmission category															
Male-to-male sexual contact	2,039	2,946	—	2,174	3,084	—	2,086	3,067	—	2,116	3,144	—	2,106	3,258	—
Injection drug use	435	687	—	347	574	—	300	522	—	272	488	—	233	463	—
Male-to-male sexual contact and injection drug use	133	182	—	127	173	—	109	156	—	98	146	—	98	147	—
Heterosexual contact ^c	660	1,088	—	646	1,035	—	603	976	—	572	973	—	484	875	—
Other ^d	1,521	8	—	1,411	10	—	1,414	15	—	1,400	7	—	1,366	8	—
Subtotal, males	4,788	4,911	17.1	4,705	4,876	16.6	4,512	4,736	15.9	4,458	4,759	15.8	4,287	4,752	15.6
Females															
Race/ethnicity															
American Indian/Alaska Native	10	10	4.9	8	8	3.8	5	5	2.4	13	14	6.2	10	11	4.7
Asian	13	13	1.0	21	22	1.5	16	17	1.1	16	17	1.1	23	26	1.6
Black/African American	1,217	1,251	35.5	1,138	1,182	32.4	1,116	1,175	31.4	1,076	1,154	30.3	975	1,090	28.0
Hispanic/Latino ^b	267	274	9.3	217	224	7.2	210	220	6.8	253	269	7.9	241	268	7.5
Native Hawaiian/Other Pacific Islander	0	0	0.0	2	2	5.5	1	1	2.5	4	4	9.9	2	2	4.9
White	299	306	1.4	296	306	1.4	286	298	1.3	304	324	1.4	279	308	1.4
Multiple races	57	59	21.0	48	50	17.1	48	50	16.6	45	49	15.6	20	22	6.7
Transmission category															
Injection drug use	209	370	—	165	313	—	165	304	—	157	291	—	111	254	—
Heterosexual contact ^c	839	1,539	—	806	1,477	—	749	1,457	—	801	1,537	—	650	1,468	—
Other ^d	815	5	—	759	4	—	768	5	—	753	3	—	789	4	—
Subtotal, females	1,863	1,913	6.3	1,730	1,795	5.7	1,682	1,766	5.6	1,711	1,831	5.7	1,550	1,725	5.4
Both sexes															
Race/ethnicity															
American Indian/Alaska Native	26	27	6.7	22	23	5.5	18	19	4.5	25	26	6.2	26	28	6.5
Asian	71	73	2.9	80	83	3.1	75	78	2.8	66	71	2.5	88	98	3.3
Black/African American	3,206	3,295	50.4	3,002	3,120	46.1	2,845	2,999	43.3	2,653	2,849	40.3	2,465	2,758	38.2
Hispanic/Latino ^b	969	993	17.3	945	976	16.1	936	981	15.4	1,010	1,076	16.1	970	1,072	15.3
Native Hawaiian/Other Pacific Islander	2	2	2.8	5	5	7.0	7	7	9.3	6	6	7.6	5	6	6.5
White	2,157	2,209	5.1	2,202	2,277	5.2	2,160	2,257	5.1	2,262	2,405	5.4	2,188	2,409	5.4
Multiple races	220	227	42.4	179	186	33.2	153	160	27.8	147	156	26.1	95	106	17.3
Total, 50–64 years of age	6,651	6,825	11.5	6,435	6,670	11.0	6,194	6,502	10.6	6,169	6,590	10.6	5,837	6,477	10.3

Table 2a. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States (cont)

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
≥65 years of age at diagnosis															
Males															
Race/ethnicity															
American Indian/Alaska Native	4	4	5.0	2	2	2.4	1	1	1.1	3	3	3.2	1	1	1.0
Asian	6	6	1.0	7	7	1.1	9	9	1.4	6	6	0.8	14	16	1.9
Black/African American	217	222	16.6	243	253	18.3	241	253	17.4	238	256	16.7	242	271	16.8
Hispanic/Latino ^b	87	89	7.5	102	106	8.3	91	96	7.1	85	91	6.3	83	91	6.0
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	1	1	6.2	0	0	0.0	1	1	5.3
White	247	253	1.8	245	254	1.8	251	264	1.7	297	317	2.0	262	291	1.8
Multiple races	18	19	16.6	16	17	14.1	10	10	8.2	16	17	12.8	8	9	6.4
Transmission category															
Male-to-male sexual contact	197	311	—	237	347	—	224	347	—	240	384	—	232	396	—
Injection drug use	25	72	—	29	69	—	22	71	—	19	65	—	24	74	—
Male-to-male sexual contact and injection drug use	7	12	—	8	13	—	4	9	—	10	16	—	3	8	—
Heterosexual contact ^c	103	186	—	118	204	—	99	198	—	101	215	—	88	193	—
Other ^d	247	13	—	223	6	—	255	9	—	275	10	—	264	8	—
Subtotal, males	579	594	3.4	615	638	3.6	604	634	3.4	645	691	3.5	611	680	3.3
Females															
Race/ethnicity															
American Indian/Alaska Native	2	2	2.0	0	0	0.0	1	1	0.9	2	2	1.9	0	0	0.0
Asian	1	1	0.1	3	3	0.4	0	0	0.0	3	3	0.3	0	0	0.0
Black/African American	132	135	6.5	131	136	6.4	133	141	6.3	139	150	6.5	143	160	6.7
Hispanic/Latino ^b	35	36	2.2	34	36	2.1	47	49	2.7	39	42	2.2	37	41	2.0
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	29	30	0.2	25	26	0.1	27	28	0.1	33	36	0.2	26	30	0.1
Multiple races	12	12	8.4	8	8	5.5	7	7	4.6	6	6	3.8	2	2	1.2
Transmission category															
Injection drug use	6	29	—	9	31	—	5	31	—	4	30	—	10	34	—
Heterosexual contact ^c	94	184	—	87	176	—	87	193	—	97	206	—	81	199	—
Other ^d	111	3	—	105	3	—	123	2	—	121	3	—	117	1	—
Subtotal, females	211	216	0.9	201	210	0.9	215	227	0.9	222	239	1.0	208	233	0.9
Both sexes															
Race/ethnicity															
American Indian/Alaska Native	6	6	3.3	2	2	1.1	2	2	1.0	5	5	2.4	1	1	0.5
Asian	7	7	0.5	10	10	0.7	9	9	0.6	9	9	0.5	14	16	0.8
Black/African American	349	358	10.5	374	389	11.1	374	394	10.7	377	406	10.6	385	431	10.7
Hispanic/Latino ^b	122	125	4.4	136	142	4.8	138	145	4.6	124	133	4.0	120	133	3.7
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	1	1	2.8	0	0	0.0	1	1	2.5
White	276	283	0.9	270	280	0.9	278	292	0.9	330	352	1.0	288	321	0.9
Multiple races	30	31	12.0	24	25	9.2	17	18	6.1	22	24	7.7	10	11	3.5
Total, ≥65 years of age at diagnosis	790	810	2.0	816	848	2.0	819	861	2.0	867	930	2.1	819	914	2.0
Total^e	7,441	7,635	7.7	7,251	7,518	7.4	7,013	7,363	7.1	7,036	7,520	7.1	6,656	7,391	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2b. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States and 6 dependent areas

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
50–64 years of age at diagnosis															
Males															
Race/ethnicity															
American Indian/Alaska Native	16	16	—	14	14	—	13	14	—	12	13	—	16	18	—
Asian	58	60	—	59	61	—	59	62	—	50	54	—	66	73	—
Black/African American	1,991	2,046	—	1,869	1,943	—	1,729	1,824	—	1,580	1,698	—	1,495	1,673	—
Hispanic/Latino ^b	804	825	—	843	873	—	810	850	—	861	921	—	808	897	—
Native Hawaiian/Other Pacific Islander	2	2	—	3	3	—	6	6	—	2	2	—	3	3	—
White	1,858	1,903	—	1,906	1,971	—	1,874	1,959	—	1,958	2,082	—	1,910	2,103	—
Multiple races	163	168	—	131	136	—	105	110	—	102	107	—	76	86	—
Transmission category															
Male-to-male sexual contact	2,064	2,974	—	2,207	3,122	—	2,112	3,096	—	2,142	3,175	—	2,133	3,293	—
Injection drug use	459	717	—	371	606	—	316	541	—	296	519	—	246	486	—
Male-to-male sexual contact and injection drug use	136	186	—	130	176	—	112	159	—	103	152	—	99	149	—
Heterosexual contact ^c	701	1,135	—	692	1,088	—	639	1,015	—	615	1,024	—	515	917	—
Other ^d	1,532	8	—	1,425	10	—	1,417	15	—	1,409	7	—	1,381	8	—
Subtotal, males	4,892	5,019	17.3	4,825	5,002	16.8	4,596	4,826	16.1	4,565	4,876	16.0	4,374	4,853	15.8
Females															
Race/ethnicity															
American Indian/Alaska Native	10	10	—	8	8	—	5	5	—	13	14	—	10	11	—
Asian	13	13	—	21	22	—	16	17	—	16	17	—	23	26	—
Black/African American	1,218	1,252	—	1,138	1,182	—	1,116	1,175	—	1,078	1,157	—	977	1,092	—
Hispanic/Latino ^b	314	323	—	258	268	—	259	273	—	304	325	—	265	296	—
Native Hawaiian/Other Pacific Islander	0	0	—	2	2	—	1	1	—	4	4	—	2	2	—
White	299	306	—	296	306	—	286	298	—	305	325	—	279	308	—
Multiple races	57	59	—	48	50	—	48	50	—	45	49	—	20	22	—
Transmission category															
Injection drug use	211	372	—	169	318	—	170	310	—	165	301	—	113	257	—
Heterosexual contact ^c	882	1,587	—	840	1,516	—	789	1,503	—	837	1,586	—	672	1,495	—
Other ^d	818	5	—	762	4	—	772	5	—	763	3	—	791	4	—
Subtotal, females	1,911	1,963	6.4	1,771	1,838	5.8	1,731	1,819	5.7	1,765	1,890	5.9	1,576	1,756	5.4
Both sexes															
Race/ethnicity															
American Indian/Alaska Native	26	27	—	22	23	—	18	19	—	25	26	—	26	28	—
Asian	71	73	—	80	83	—	75	78	—	66	71	—	89	99	—
Black/African American	3,209	3,298	—	3,007	3,125	—	2,845	2,999	—	2,658	2,855	—	2,472	2,766	—
Hispanic/Latino ^b	1,118	1,148	—	1,101	1,141	—	1,069	1,123	—	1,165	1,246	—	1,073	1,192	—
Native Hawaiian/Other Pacific Islander	2	2	—	5	5	—	7	7	—	6	6	—	5	6	—
White	2,157	2,209	—	2,202	2,277	—	2,160	2,257	—	2,263	2,406	—	2,189	2,411	—
Multiple races	220	227	—	179	186	—	153	160	—	147	156	—	96	107	—
Total, 50–64 years of age	6,803	6,983	11.7	6,596	6,840	11.1	6,327	6,645	10.7	6,330	6,766	10.8	5,950	6,608	10.4

Table 2b. Diagnoses of HIV infection among adults aged 50 years and older, by year of diagnosis, age group, and selected characteristics, 2010–2014—United States and 6 dependent areas (cont)

	2010			2011			2012			2013			2014		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate												
≥65 years of age at diagnosis															
Males															
Race/ethnicity															
American Indian/Alaska Native	4	4	—	2	2	—	1	1	—	3	3	—	1	1	—
Asian	6	6	—	7	7	—	9	9	—	6	6	—	14	16	—
Black/African American	218	223	—	243	253	—	241	253	—	238	256	—	243	272	—
Hispanic/Latino ^b	108	111	—	115	120	—	115	121	—	106	114	—	94	104	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	1	1	—	0	0	—	1	1	—
White	247	253	—	245	254	—	251	264	—	297	317	—	262	291	—
Multiple races	18	19	—	16	17	—	10	10	—	16	17	—	8	9	—
Transmission category															
Male-to-male sexual contact	200	314	—	242	352	—	228	352	—	243	388	—	235	400	—
Injection drug use	29	77	—	29	69	—	27	77	—	23	70	—	27	78	—
Male-to-male sexual contact and injection drug use	7	12	—	8	13	—	4	9	—	10	16	—	3	8	—
Heterosexual contact ^c	115	201	—	126	212	—	113	214	—	114	230	—	94	200	—
Other ^d	250	13	—	223	6	—	256	9	—	276	10	—	264	8	—
Subtotal, males	601	617	3.5	628	652	3.6	628	660	3.5	666	714	3.6	623	694	3.4
Females															
Race/ethnicity															
American Indian/Alaska Native	2	2	—	0	0	—	1	1	—	2	2	—	0	0	—
Asian	1	1	—	3	3	—	0	0	—	3	3	—	0	0	—
Black/African American	132	135	—	131	136	—	133	141	—	139	150	—	143	160	—
Hispanic/Latino ^b	42	43	—	46	48	—	61	64	—	44	47	—	41	46	—
Native Hawaiian/Other Pacific Islander	29	30	—	25	26	—	27	28	—	33	36	—	26	30	—
White	12	12	—	8	8	—	7	7	—	6	6	—	2	2	—
Multiple races			—			—			—			—			—
Transmission category															
Injection drug use	6	29	—	9	32	—	5	31	—	5	31	—	10	34	—
Heterosexual contact ^c	101	192	—	98	188	—	101	208	—	100	211	—	85	203	—
Other ^d	111	3	—	106	3	—	123	2	—	122	3	—	117	1	—
Subtotal, females	218	224	1.0	213	222	0.9	229	242	1.0	227	245	1.0	212	238	0.9
Both sexes															
Race/ethnicity															
American Indian/Alaska Native	6	6	—	2	2	—	2	2	—	5	5	—	1	1	—
Asian	7	7	—	10	10	—	9	9	—	9	9	—	14	16	—
Black/African American	350	359	—	374	389	—	374	394	—	377	406	—	386	432	—
Hispanic/Latino ^b	150	155	—	161	168	—	176	186	—	150	162	—	135	150	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	1	1	—	0	0	—	1	1	—
White	276	283	—	270	280	—	278	292	—	330	352	—	288	321	—
Multiple races	30	31	—	24	25	—	17	18	—	22	24	—	10	11	—
Total, ≥65 years of age at diagnosis	819	840	2.0	841	874	2.1	857	902	2.1	893	959	2.1	835	933	2.0
Total^e	7,622	7,823	7.8	7,437	7,714	7.5	7,184	7,546	7.1	7,223	7,725	7.2	6,785	7,541	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3a. Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States

	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	13	14	17.8	3	4	5.1	0	0	0.0	0	0	0.0
Asian	43	48	9.2	15	17	3.6	7	8	2.0	11	12	4.2
Black/African American	693	773	60.5	531	594	51.5	266	301	33.8	154	173	27.3
Hispanic/Latino ^b	404	446	30.4	214	237	20.9	111	121	14.8	48	53	9.1
Native Hawaiian/Other Pacific Islander	1	1	6.4	0	0	0.0	2	2	21.3	0	0	0.0
White	1,030	1,133	14.9	580	641	8.5	299	328	4.9	160	177	3.1
Multiple races	44	49	43.2	26	29	28.8	5	6	7.8	4	4	7.9
Transmission category												
Male-to-male sexual contact	1,175	1,772	—	623	1,004	—	308	483	—	147	253	—
Injection drug use	102	198	—	83	173	—	48	92	—	18	47	—
Male-to-male sexual contact and injection drug use	51	77	—	32	48	—	15	22	—	3	6	—
Heterosexual contact ^c	254	415	—	150	294	—	80	166	—	49	110	—
Other ^d	646	3	—	481	2	—	239	3	—	160	3	—
Subtotal, males	2,228	2,464	22.2	1,369	1,521	14.6	690	766	8.6	377	420	5.8
Females												
Race/ethnicity												
American Indian/Alaska Native	6	6	7.5	3	3	4.1	1	1	1.7	0	0	0.0
Asian	9	10	1.7	10	11	2.1	4	4	0.9	0	0	0.0
Black/African American	421	470	32.4	377	420	31.3	177	200	18.2	100	112	13.7
Hispanic/Latino ^b	111	123	8.3	83	93	7.8	47	52	5.7	18	20	2.9
Native Hawaiian/Other Pacific Islander	1	1	6.3	1	1	7.2	0	0	0.0	0	0	0.0
White	141	156	2.0	102	112	1.4	36	40	0.6	17	20	0.3
Multiple races	11	12	9.5	4	4	4.0	5	5	6.2	1	1	1.6
Transmission category												
Injection drug use	53	114	—	40	95	—	18	45	—	8	23	—
Heterosexual contact ^c	301	662	—	241	549	—	108	257	—	51	130	—
Other ^d	346	2	—	299	1	—	144	1	—	77	0	—
Subtotal, females	700	778	6.8	580	645	5.8	270	303	3.1	136	153	1.9
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	19	20	12.5	6	7	4.6	1	1	0.9	0	0	0.0
Asian	52	58	5.2	25	28	2.8	11	12	1.4	11	12	1.9
Black/African American	1,114	1,242	45.6	908	1,014	40.6	443	501	25.2	254	285	19.7
Hispanic/Latino ^b	515	569	19.4	297	330	14.2	158	173	10.0	66	73	5.7
Native Hawaiian/Other Pacific Islander	2	2	6.3	1	1	3.6	2	2	10.4	0	0	0.0
White	1,171	1,289	8.4	682	753	4.9	335	368	2.7	177	197	1.7
Multiple races	55	61	25.5	30	33	15.8	10	11	7.0	5	6	4.5
Total^e	2,928	3,242	14.4	1,949	2,166	10.1	960	1,069	5.8	513	573	3.7

Table 3a. Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States (cont)

	70–74 years			75–79 years			80–84 years			≥85 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	1	1	3.8	0	0	0.0	0	0	0.0	0	0	0.0
Asian	3	3	1.6	0	0	0.0	0	0	0.0	0	0	0.0
Black/African American	49	54	13.2	25	28	10.2	12	14	8.5	2	2	1.6
Hispanic/Latino ^b	22	24	6.4	7	8	3.0	6	7	3.9	0	0	0.0
Native Hawaiian/Other Pacific Islander	1	1	20.3	0	0	0.0	0	0	0.0	0	0	0.0
White	58	65	1.6	39	43	1.6	4	4	0.2	1	1	0.1
Multiple races	1	1	2.9	2	3	10.6	1	1	8.2	0	0	0.0
Transmission category												
Male-to-male sexual contact	50	84	—	29	49	—	6	11	—	0	0	—
Injection drug use	4	15	—	2	9	—	0	3	—	0	0	—
Male-to-male sexual contact and injection drug use	0	1	—	0	0	—	0	0	—	0	0	—
Heterosexual contact ^c	26	48	—	9	21	—	4	12	—	0	2	—
Other ^d	55	2	—	33	2	—	13	1	—	3	1	—
Subtotal, males	135	150	2.9	73	81	2.3	23	26	1.1	3	3	0.1
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
Asian	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
Black/African American	31	35	6.1	6	6	1.5	2	2	0.7	4	5	1.5
Hispanic/Latino ^b	12	13	2.7	6	7	1.9	1	1	0.6	0	0	0.0
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	5	5	0.1	3	3	0.1	0	0	0.0	1	1	0.0
Multiple races	1	1	2.4	0	0	0.0	0	0	0.0	0	0	0.0
Transmission category												
Injection drug use	1	7	—	1	2	—	0	0	—	0	1	—
Heterosexual contact ^c	18	47	—	9	14	—	1	3	—	2	5	—
Other ^d	30	1	—	5	0	—	2	0	—	3	0	—
Subtotal, females	49	55	0.9	15	17	0.4	3	4	0.1	5	6	0.1
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	1	1	1.8	0	0	0.0	0	0	0.0	0	0	0.0
Asian	3	3	0.7	0	0	0.0	0	0	0.0	0	0	0.0
Black/African American	80	89	9.1	31	34	5.0	14	16	3.5	6	7	1.5
Hispanic/Latino ^b	34	37	4.3	13	15	2.4	7	8	1.9	0	0	0.0
Native Hawaiian/Other Pacific Islander	1	1	9.8	0	0	0.0	0	0	0.0	0	0	0.0
White	63	71	0.8	42	47	0.8	4	4	0.1	2	2	0.0
Multiple races	2	2	2.6	2	3	4.7	1	1	3.3	0	0	0.0
Total^e	184	205	1.9	88	98	1.2	26	30	0.5	8	9	0.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3b. Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States and 6 dependent areas

	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	13	14	—	3	4	—	0	0	—	0	0	—
Asian	44	49	—	15	17	—	7	8	—	11	12	—
Black/African American	696	776	—	532	595	—	267	302	—	155	174	—
Hispanic/Latino ^b	433	480	—	243	271	—	132	145	—	53	59	—
Native Hawaiian/Other Pacific Islander	1	1	—	0	0	—	2	2	—	0	0	—
White	1,031	1,134	—	580	641	—	299	328	—	160	177	—
Multiple races	44	49	—	27	30	—	5	6	—	4	4	—
Transmission category												
Male-to-male sexual contact	1,189	1,790	—	630	1,013	—	314	490	—	148	254	—
Injection drug use	105	204	—	90	183	—	51	98	—	21	50	—
Male-to-male sexual contact and injection drug use	51	77	—	33	49	—	15	22	—	3	6	—
Heterosexual contact ^c	265	430	—	162	309	—	88	178	—	51	113	—
Other ^d	652	3	—	485	2	—	244	3	—	160	3	—
Subtotal, males	2,262	2,504	22.4	1,400	1,558	14.8	712	792	8.8	383	427	5.8
Females												
Race/ethnicity												
American Indian/Alaska Native	6	6	—	3	3	—	1	1	—	0	0	—
Asian	9	10	—	10	11	—	4	4	—	0	0	—
Black/African American	422	471	—	378	422	—	177	200	—	100	112	—
Hispanic/Latino ^b	121	134	—	93	104	—	51	57	—	19	21	—
Native Hawaiian/Other Pacific Islander	1	1	—	1	1	—	0	0	—	0	0	—
White	141	156	—	102	112	—	36	40	—	17	20	—
Multiple races	11	12	—	4	4	—	5	5	—	1	1	—
Transmission category												
Injection drug use	54	115	—	41	97	—	18	45	—	8	23	—
Heterosexual contact ^c	310	674	—	250	560	—	112	261	—	52	131	—
Other ^d	347	2	—	300	1	—	144	1	—	77	0	—
Subtotal, females	711	791	6.8	591	658	5.9	274	307	3.1	137	154	1.9
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	19	20	—	6	7	—	1	1	—	0	0	—
Asian	53	59	—	25	28	—	11	12	—	11	12	—
Black/African American	1,118	1,247	—	910	1,016	—	444	502	—	255	286	—
Hispanic/Latino ^b	554	614	—	336	376	—	183	202	—	72	79	—
Native Hawaiian/Other Pacific Islander	2	2	—	1	1	—	2	2	—	0	0	—
White	1,172	1,290	—	682	753	—	335	368	—	177	197	—
Multiple races	55	61	—	31	35	—	10	11	—	5	6	—
Total^e	2,973	3,294	14.4	1,991	2,215	10.2	986	1,099	5.8	520	581	3.7

Table 3b. Diagnoses of HIV infection among adults aged 50 years and older, by 5-year age group and selected characteristics, 2014—United States and 6 dependent areas (cont)

	70–74 years			75–79 years			80–84 years			≥85 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	1	1	—	0	0	—	0	0	—	0	0	—
Asian	3	3	—	0	0	—	0	0	—	0	0	—
Black/African American	49	54	—	25	28	—	12	14	—	2	2	—
Hispanic/Latino ^b	26	29	—	9	10	—	6	7	—	0	0	—
Native Hawaiian/Other Pacific Islander	1	1	—	0	0	—	0	0	—	0	0	—
White	58	65	—	39	43	—	4	4	—	1	1	—
Multiple races	1	1	—	2	3	—	1	1	—	0	0	—
Transmission category												
Male-to-male sexual contact	51	85	—	30	50	—	6	11	—	0	0	—
Injection drug use	4	15	—	2	9	—	0	3	—	0	0	—
Male-to-male sexual contact and injection drug use	0	1	—	0	0	—	0	0	—	0	0	—
Heterosexual contact ^c	29	52	—	10	22	—	4	12	—	0	2	—
Other ^d	55	2	—	33	2	—	13	1	—	3	1	—
Subtotal, males	139	155	3.0	75	84	2.3	23	26	1.1	3	3	0.1
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	—	0	0	—	0	0	—	0	0	—
Asian	0	0	—	0	0	—	0	0	—	0	0	—
Black/African American	31	35	—	6	6	—	2	2	—	4	5	—
Hispanic/Latino ^b	15	17	—	6	7	—	1	1	—	0	0	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	0	0	—
White	5	5	—	3	3	—	0	0	—	1	1	—
Multiple races	1	1	—	0	0	—	0	0	—	0	0	—
Transmission category												
Injection drug use	1	7	—	1	2	—	0	0	—	0	1	—
Heterosexual contact ^c	21	50	—	9	14	—	1	3	—	2	5	—
Other ^d	30	1	—	5	0	—	2	0	—	3	0	—
Subtotal, females	52	58	1.0	15	17	0.4	3	4	0.1	5	6	0.1
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	1	1	—	0	0	—	0	0	—	0	0	—
Asian	3	3	—	0	0	—	0	0	—	0	0	—
Black/African American	80	89	—	31	34	—	14	16	—	6	7	—
Hispanic/Latino ^b	41	46	—	15	17	—	7	8	—	0	0	—
Native Hawaiian/Other Pacific Islander	1	1	—	0	0	—	0	0	—	0	0	—
White	63	71	—	42	47	—	4	4	—	2	2	—
Multiple races	2	2	—	2	3	—	1	1	—	0	0	—
Total^e	191	213	1.9	90	100	1.2	26	30	0.5	8	9	0.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4a. Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States

	American Indian/ Alaska Native			Asian			Black/African American			Hispanic/Latino ^a		
	No.	Estimated ^b		No.	Estimated ^b		No.	Estimated ^b		No.	Estimated ^b	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
50–64 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	8	10	-	33	58	-	406	798	-	363	559	-
Injection drug use	3	4	-	1	2	-	93	233	-	50	101	-
Male-to-male sexual contact and injection drug use	2	2	-	0	0	-	23	42	-	11	20	-
Heterosexual contact ^d	1	2	-	5	11	-	303	590	-	75	124	-
Other ^e	2	0	-	26	0	-	665	4	-	230	1	-
Subtotal, males	16	18	8.6	65	72	5.3	1,490	1,668	50.2	729	805	23.5
Females												
Transmission category												
Injection drug use	2	3	-	1	3	-	54	142	-	19	44	-
Heterosexual contact ^d	6	8	-	7	22	-	399	947	-	104	223	-
Other ^e	2	0	-	15	1	-	522	1	-	118	0	-
Subtotal, females	10	11	4.7	23	26	1.6	975	1,090	28.0	241	268	7.5
Subtotal, 50–64 years of age at diagnosis	26	28	6.5	88	98	3.3	2,465	2,758	38.2	970	1,072	15.3
≥65 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	0	1	-	2	9	-	47	91	-	37	56	-
Injection drug use	0	0	-	0	1	-	19	48	-	2	10	-
Male-to-male sexual contact and injection drug use	0	0	-	0	0	-	2	4	-	1	2	-
Heterosexual contact ^d	0	0	-	1	5	-	54	126	-	13	23	-
Other ^e	1	0	-	11	1	-	120	2	-	30	1	-
Subtotal, males	1	1	1.0	14	16	1.9	242	271	16.8	83	91	6.0
Females												
Transmission category												
Injection drug use	0	0	-	0	0	-	5	20	-	2	7	-
Heterosexual contact ^d	0	0	-	0	0	-	57	140	-	15	34	-
Other ^e	0	0	-	0	0	-	81	0	-	20	0	-
Subtotal, females	0	0	0.0	0	0	0.0	143	160	6.7	37	41	2.0
Subtotal, ≥65 years of age at diagnosis	1	1	0.5	14	16	0.8	385	431	10.7	120	133	3.7
Total, ≥50 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	8	11	-	35	67	-	453	889	-	400	615	-
Injection drug use	3	4	-	1	3	-	112	282	-	52	111	-
Male-to-male sexual contact and injection drug use	2	2	-	0	0	-	25	46	-	12	22	-
Heterosexual contact ^d	1	2	-	6	16	-	357	715	-	88	148	-
Other ^e	3	0	-	37	1	-	785	7	-	260	1	-
Subtotal, males	17	19	6.0	79	88	4.1	1,732	1,938	39.3	812	896	18.1
Females												
Transmission category												
Injection drug use	2	3	-	1	3	-	59	162	-	21	51	-
Heterosexual contact ^d	6	8	-	7	22	-	456	1,087	-	119	257	-
Other ^e	2	0	-	15	1	-	603	1	-	138	1	-
Subtotal, females	10	11	3.0	23	26	1.0	1,118	1,250	19.9	278	309	5.5
Total^f	27	29	4.4	102	114	2.3	2,850	3,189	28.4	1,090	1,205	11.4

Table 4a. Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States (cont)

	Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^c	Rate
50–64 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	1	2	-	1,259	1,766	-	36	64	-	2,106	3,258	-
Injection drug use	1	1	-	81	114	-	4	7	-	233	463	-
Male-to-male sexual contact and injection drug use	0	0	-	61	80	-	1	3	-	98	147	-
Heterosexual contact ^d	0	0	-	94	139	-	6	10	-	484	875	-
Other ^e	1	0	-	414	3	-	28	0	-	1,366	8	-
Subtotal, males	3	3	8.1	1,909	2,102	9.7	75	85	28.9	4,287	4,752	15.6
Females												
Transmission category												
Injection drug use	0	0	-	31	56	-	4	7	-	111	254	-
Heterosexual contact ^d	1	2	-	128	251	-	5	15	-	650	1,468	-
Other ^e	1	0	-	120	1	-	11	0	-	789	4	-
Subtotal, females	2	2	4.9	279	308	1.4	20	22	6.7	1,550	1,725	5.4
Subtotal, 50–64 years of age at diagnosis	5	6	6.5	2,188	2,409	5.4	95	106	17.3	5,837	6,477	10.3
≥65 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	1	1	-	143	234	-	2	5	-	232	396	-
Injection drug use	0	0	-	3	14	-	0	1	-	24	74	-
Male-to-male sexual contact and injection drug use	0	0	-	0	3	-	0	0	-	3	8	-
Heterosexual contact ^d	0	0	-	19	36	-	1	3	-	88	193	-
Other ^e	0	0	-	97	4	-	5	0	-	264	8	-
Subtotal, males	1	1	5.3	262	291	1.8	8	9	6.4	611	680	3.3
Females												
Transmission category												
Injection drug use	0	0	-	2	6	-	1	1	-	10	34	-
Heterosexual contact ^d	0	0	-	8	24	-	1	1	-	81	199	-
Other ^e	0	0	-	16	0	-	0	0	-	117	1	-
Subtotal, females	0	0	0.0	26	30	0.1	2	2	1.2	208	233	0.9
Subtotal, ≥65 years of age at diagnosis	1	1	2.5	288	321	0.9	10	11	3.5	819	914	2.0
Total, ≥50 years of age at diagnosis												
Males												
Transmission category												
Male-to-male sexual contact	2	3	-	1,402	2,000	-	38	69	-	2,338	3,654	-
Injection drug use	1	1	-	84	129	-	4	8	-	257	537	-
Male-to-male sexual contact and injection drug use	0	0	-	61	83	-	1	3	-	101	155	-
Heterosexual contact ^d	0	0	-	113	175	-	7	13	-	572	1,069	-
Other ^e	1	0	-	511	7	-	33	0	-	1,630	17	-
Subtotal, males	4	4	7.2	2,171	2,393	6.3	83	94	21.5	4,898	5,432	10.7
Females												
Transmission category												
Injection drug use	0	0	-	33	62	-	5	8	-	121	288	-
Heterosexual contact ^d	1	2	-	136	274	-	6	16	-	731	1,667	-
Other ^e	1	0	-	136	2	-	11	0	-	906	4	-
Subtotal, females	2	2	3.2	305	337	0.8	22	24	4.7	1,758	1,959	3.4
Total^f	6	7	5.1	2,476	2,730	3.4	105	118	12.5	6,656	7,391	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4b. Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian		Black/African American		Hispanic/Latino ^a	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b
50–64 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	8	10	34	59	406	798	387	591
Injection drug use	3	4	1	2	93	235	63	121
Male-to-male sexual contact and injection drug use	2	2	0	0	23	42	12	21
Heterosexual contact ^d	1	2	5	11	305	593	104	162
Other ^e	2	0	26	0	668	4	242	1
Subtotal, males	16	18	66	73	1,495	1,673	808	897
Females								
Transmission category								
Injection drug use	2	3	1	3	54	142	21	47
Heterosexual contact ^d	6	8	7	22	400	949	125	249
Other ^e	2	0	15	1	523	1	119	0
Subtotal, females	10	11	23	26	977	1,092	265	296
Subtotal, 50–64 years of age at diagnosis	26	28	89	99	2,472	2,766	1,073	1,192
≥65 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	0	1	2	9	47	91	40	59
Injection drug use	0	0	0	1	19	48	5	13
Male-to-male sexual contact and injection drug use	0	0	0	0	2	4	1	2
Heterosexual contact ^d	0	0	1	5	55	127	18	29
Other ^e	1	0	11	1	120	2	30	1
Subtotal, males	1	1	14	16	243	272	94	104
Females								
Transmission category								
Injection drug use	0	0	0	0	5	20	2	7
Heterosexual contact ^d	0	0	0	0	57	140	19	39
Other ^e	0	0	0	0	81	0	20	0
Subtotal, females	0	0	0	0	143	160	41	46
Subtotal, ≥65 years of age at diagnosis	1	1	14	16	386	432	135	150
Total, ≥50 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	8	11	36	68	453	889	427	650
Injection drug use	3	4	1	3	112	283	68	135
Male-to-male sexual contact and injection drug use	2	2	0	0	25	46	13	23
Heterosexual contact ^d	1	2	6	16	360	720	122	192
Other ^e	3	0	37	1	788	7	272	1
Subtotal, males	17	19	80	89	1,738	1,945	902	1,001
Females								
Transmission category								
Injection drug use	2	3	1	3	59	162	23	54
Heterosexual contact ^d	6	8	7	22	457	1,089	144	287
Other ^e	2	0	15	1	604	1	139	1
Subtotal, females	10	11	23	26	1,120	1,252	306	341
Total^f	27	29	103	115	2,858	3,198	1,208	1,342

Table 4b. Diagnoses of HIV infection among adults aged 50 years and older, by race/ethnicity, age group, and selected characteristics, 2014—United States and 6 dependent areas (cont)

	Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^{b,c}
50–64 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	1	2	1,260	1,767	37	66	2,133	3,293
Injection drug use	1	1	81	114	4	7	246	486
Male-to-male sexual contact and injection drug use	0	0	61	80	1	3	99	149
Heterosexual contact ^d	0	0	94	139	6	10	515	917
Other ^e	1	0	414	3	28	0	1,381	8
Subtotal, males	3	3	1,910	2,103	76	86	4,374	4,853
Females								
Transmission category								
Injection drug use	0	0	31	56	4	7	113	257
Heterosexual contact ^d	1	2	128	251	5	15	672	1,495
Other ^e	1	0	120	1	11	0	791	4
Subtotal, females	2	2	279	308	20	22	1,576	1,756
Subtotal, 50–64 years of age at diagnosis	5	6	2,189	2,411	96	107	5,950	6,608
≥65 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	1	1	143	234	2	5	235	400
Injection drug use	0	0	3	14	0	1	27	78
Male-to-male sexual contact and injection drug use	0	0	0	3	0	0	3	8
Heterosexual contact ^d	0	0	19	36	1	3	94	200
Other ^e	0	0	97	4	5	0	264	8
Subtotal, males	1	1	262	291	8	9	623	694
Females								
Transmission category								
Injection drug use	0	0	2	6	1	1	10	34
Heterosexual contact ^d	0	0	8	24	1	1	85	203
Other ^e	0	0	16	0	0	0	117	1
Subtotal, females	0	0	26	30	2	2	212	238
Subtotal, ≥65 years of age at diagnosis	1	1	288	321	10	11	835	933
Total, ≥50 years of age at diagnosis								
Males								
Transmission category								
Male-to-male sexual contact	2	3	1,403	2,001	39	70	2,368	3,693
Injection drug use	1	1	84	129	4	8	273	563
Male-to-male sexual contact and injection drug use	0	0	61	83	1	3	102	157
Heterosexual contact ^d	0	0	113	175	7	13	609	1,117
Other ^e	1	0	511	7	33	0	1,645	17
Subtotal, males	4	4	2,172	2,394	84	95	4,997	5,547
Females								
Transmission category								
Injection drug use	0	0	33	62	5	8	123	291
Heterosexual contact ^d	1	2	136	274	6	16	757	1,699
Other ^e	1	0	136	2	11	0	908	4
Subtotal, females	2	2	305	337	22	24	1,788	1,994
Total^f	6	7	2,477	2,731	106	119	6,785	7,541

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 5. Diagnoses of HIV infection among adults aged 50 years and older, by age group and area of residence, 2014—United States and 6 dependent areas

Area of residence	Age at diagnosis						Total		
	50–64 years			≥65 years					
	No. ^b	Estimated ^a		No. ^b	Estimated ^a		No. ^b	Estimated ^a	
		No.	Rate		No.	Rate		No. ^c	Rate
Alabama	67	74	7.6	12	13	1.8	79	88	5.1
Alaska	6	6	4.4	0	0	0.0	6	6	3.0
Arizona	95	101	8.2	14	15	1.4	109	116	5.0
Arkansas	41	45	7.9	4	5	1.0	45	50	4.8
California	681	773	10.8	81	93	1.9	762	866	7.1
Colorado	37	39	3.7	5	5	0.8	42	44	2.6
Connecticut	61	65	8.4	12	13	2.3	73	77	5.8
Delaware	18	21	10.7	3	3	2.3	21	24	6.9
District of Columbia	55	64	59.7	10	12	15.6	65	76	41.7
Florida	985	1,045	26.1	148	157	4.1	1,133	1,202	15.4
Georgia	230	315	16.7	26	36	2.9	256	351	11.2
Hawaii	16	17	6.2	3	3	1.4	19	20	4.0
Idaho	6	7	2.2	0	0	0.0	6	7	1.3
Illinois	172	197	7.8	26	29	1.6	198	227	5.2
Indiana	66	71	5.4	4	4	0.4	70	75	3.3
Iowa	24	25	4.1	2	2	0.4	26	27	2.5
Kansas	16	17	3.1	1	1	0.3	17	18	1.9
Kentucky	47	50	5.6	2	2	0.3	49	52	3.4
Louisiana	177	187	20.5	20	21	3.3	197	208	13.5
Maine	15	16	5.0	5	5	2.2	20	21	3.8
Maryland	188	235	19.2	25	31	3.8	213	266	13.0
Massachusetts	102	139	10.0	17	23	2.3	119	162	6.7
Michigan	95	101	4.8	16	17	1.1	111	118	3.2
Minnesota	34	36	3.3	7	7	0.9	41	44	2.3
Mississippi	65	69	12.0	5	6	1.3	70	75	7.4
Missouri	57	60	4.9	6	6	0.7	63	67	3.1
Montana	1	1	0.5	0	0	0.0	1	1	0.3
Nebraska	10	11	2.9	1	1	0.4	11	12	1.8
Nevada	63	67	12.4	3	3	0.8	66	70	7.5
New Hampshire	9	10	3.1	1	1	0.5	10	11	2.0
New Jersey	254	296	16.0	41	49	3.8	295	345	10.9
New Mexico	23	24	5.9	1	1	0.4	24	26	3.5
New York	543	590	15.0	107	117	4.0	650	708	10.3

Table 5. Diagnoses of HIV infection among adults aged 50 years and older, by age group and area of residence, 2014—United States and 6 dependent areas (cont)

Area of residence	Age at diagnosis								
	50–64 years			≥65 years			Total		
	No. ^b	Estimated ^a		No. ^b	Estimated ^a		No. ^b	Estimated ^a	
		No.	Rate		No.	Rate		No. ^c	Rate
North Carolina	188	199	10.3	22	24	1.6	210	223	6.6
North Dakota	6	6	4.5	0	0	0.0	6	6	2.6
Ohio	95	102	4.2	12	13	0.7	107	115	2.7
Oklahoma	43	48	6.5	3	3	0.6	46	51	3.9
Oregon	36	38	4.7	5	5	0.8	41	44	3.0
Pennsylvania	225	241	8.9	32	34	1.6	257	275	5.7
Rhode Island	16	17	7.6	1	1	0.6	17	18	4.6
South Carolina	129	137	14.1	19	21	2.7	148	157	9.1
South Dakota	5	5	3.1	0	0	0.0	5	5	1.7
Tennessee	96	102	7.8	17	18	1.8	113	120	5.3
Texas	490	543	11.5	67	74	2.4	557	617	7.9
Utah	11	12	2.7	2	2	0.7	13	14	1.9
Vermont	2	2	1.5	1	1	1.2	3	3	1.3
Virginia	127	135	8.2	16	18	1.5	143	153	5.5
Washington	55	58	4.1	12	13	1.3	67	71	3.0
West Virginia	13	15	3.6	1	1	0.3	14	16	2.1
Wisconsin	41	44	3.6	1	1	0.1	42	45	2.1
Wyoming	0	0	0.0	0	0	0.0	0	0	0.0
Subtotal	5,837	6,477	10.3	819	914	2.0	6,656	7,391	6.8
U.S. dependent areas									
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0
Guam	0	0	0.0	0	0	0.0	0	0	0.0
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0
Puerto Rico	103	120	17.9	15	18	2.9	118	137	10.7
Republic of Palau	0	0	0.0	0	0	0.0	0	0	0.0
U.S. Virgin Islands	10	11	48.9	1	1	6.0	11	13	29.5
Subtotal	113	131	17.8	16	19	2.9	129	150	10.8
Total^d	5,950	6,608	10.4	835	933	2.0	6,785	7,541	6.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 6a. Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
Males												
Age at end of year (yr)												
50–54	104,318	104,302	951.6	111,297	111,257	1,004.6	118,316	118,249	1,066.4	125,610	125,447	1,131.7
55–59	67,661	67,512	704.1	74,560	74,349	757.1	81,697	81,405	808.0	88,563	88,117	856.5
60–64	36,491	36,353	445.6	41,604	41,391	484.3	46,676	46,361	542.9	52,278	51,803	597.0
65–69	15,901	15,813	268.2	18,280	18,146	298.6	21,382	21,183	320.3	24,948	24,613	356.1
70–74	6,827	6,777	158.6	7,920	7,844	178.1	9,222	9,103	197.9	10,725	10,536	215.7
75–79	2,717	2,686	84.3	3,142	3,095	95.7	3,569	3,502	106.3	4,186	4,085	120.4
80–84	951	934	40.5	1,109	1,085	46.4	1,342	1,305	55.4	1,553	1,496	63.1
≥85	379	370	20.4	462	449	23.8	537	516	26.3	657	624	30.5
Race/ethnicity												
American Indian/Alaska Native	537	534	198.1	594	590	210.9	661	656	226.1	737	730	243.1
Asian ^b	2,028	2,034	114.9	2,234	2,241	120.5	2,558	2,567	131.4	2,856	2,869	139.4
Black/African American	84,391	84,157	1,934.9	91,555	91,204	2,027.0	98,993	98,476	2,119.9	106,785	105,943	2,212.9
Hispanic/Latino ^c	39,141	39,150	983.2	43,455	43,442	1,031.2	48,170	48,129	1,081.3	53,167	53,053	1,129.6
Native Hawaiian/Other Pacific Islander	166	167	335.7	185	186	352.8	208	209	374.9	226	227	385.9
White	101,555	101,282	286.2	112,252	111,872	310.0	123,406	122,884	334.5	135,244	134,474	360.2
Multiple races	7,427	7,422	2,020.0	8,099	8,081	2,099.6	8,745	8,704	2,164.7	9,505	9,427	2,247.8
Transmission category												
Male-to-male sexual contact	121,454	138,870	—	135,605	155,160	—	150,731	172,615	—	166,864	191,083	—
Injection drug use	40,030	47,881	—	41,583	49,924	—	43,006	51,831	—	44,428	53,622	—
Male-to-male sexual contact and injection drug use	16,489	18,118	—	18,085	19,848	—	19,709	21,611	—	21,449	23,452	—
Heterosexual contact ^d	21,859	28,798	—	23,982	31,536	—	26,139	34,346	—	28,438	37,280	—
Other ^e	35,413	1,079	—	39,119	1,148	—	43,156	1,223	—	47,341	1,286	—
Subtotal, males	235,245	234,746	508.3	258,374	257,616	543.8	282,741	281,625	580.3	308,520	306,722	617.7
Females												
Age at end of year (yr)												
50–54	30,775	30,803	270.4	32,751	32,777	284.9	34,717	34,735	301.7	36,597	36,587	318.0
55–59	19,236	19,231	188.4	21,555	21,533	206.1	23,885	23,845	222.6	26,276	26,188	239.8
60–64	9,763	9,753	110.4	11,309	11,288	121.7	12,929	12,883	138.7	14,621	14,531	153.7
65–69	4,234	4,225	63.8	4,879	4,864	71.4	5,679	5,652	76.6	6,718	6,663	86.5
70–74	1,913	1,904	37.6	2,202	2,189	42.0	2,605	2,585	47.7	3,009	2,973	51.9
75–79	846	841	20.3	972	962	23.1	1,114	1,099	26.2	1,309	1,286	30.0
80–84	322	318	9.2	367	362	10.5	454	446	13.0	526	514	15.1
≥85	153	149	4.0	169	163	4.3	185	176	4.5	217	204	5.1

Table 6a. Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
Race/ethnicity												
American Indian/Alaska Native	214	213	69.5	242	241	75.4	261	260	78.3	287	286	83.3
Asian ^b	467	469	21.4	522	525	22.8	562	566	23.4	624	627	24.6
Black/African American	39,641	39,632	707.9	43,715	43,682	756.0	48,245	48,176	809.2	52,889	52,714	860.5
Hispanic/Latino ^c	11,914	11,932	261.2	13,117	13,130	272.7	14,400	14,405	284.5	15,675	15,662	294.3
Native Hawaiian/Other Pacific Islander	41	41	77.0	43	43	76.5	48	48	80.9	59	60	94.6
White	12,546	12,514	31.1	13,894	13,845	33.8	15,172	15,095	36.3	16,684	16,563	39.3
Multiple races	2,419	2,421	568.8	2,671	2,670	600.3	2,880	2,872	618.5	3,055	3,035	627.2
Transmission category												
Injection drug use	17,027	21,899	—	18,289	23,612	—	19,529	25,315	—	20,842	27,057	—
Heterosexual contact ^d	32,842	44,480	—	36,470	49,659	—	40,278	55,208	—	44,229	60,962	—
Other ^e	17,373	845	—	19,445	867	—	21,761	898	—	24,202	927	—
Subtotal, females	67,242	67,224	125.8	74,204	74,137	135.6	81,568	81,421	145.8	89,273	88,946	156.0
Both sexes												
Age at end of year (yr)												
50–54	135,093	135,105	604.4	144,048	144,034	637.9	153,033	152,984	676.9	162,207	162,034	717.2
55–59	86,897	86,743	438.2	96,115	95,882	473.0	105,582	105,251	506.3	114,839	114,306	538.9
60–64	46,254	46,106	271.4	52,913	52,679	295.6	59,605	59,243	332.3	66,899	66,334	365.8
65–69	20,135	20,038	160.0	23,159	23,010	178.6	27,061	26,835	191.8	31,666	31,276	214.0
70–74	8,740	8,681	93.0	10,122	10,033	104.4	11,827	11,688	116.7	13,734	13,509	127.3
75–79	3,563	3,526	48.2	4,114	4,057	54.9	4,683	4,602	61.4	5,495	5,372	69.9
80–84	1,273	1,253	21.8	1,476	1,447	25.0	1,796	1,751	30.3	2,079	2,009	34.8
≥85	532	518	9.4	631	611	10.7	722	692	11.8	874	828	13.7
Race/ethnicity												
American Indian/Alaska Native	751	748	129.6	836	831	138.7	922	916	147.3	1,024	1,015	157.8
Asian ^b	2,495	2,503	63.1	2,756	2,766	66.4	3,120	3,133	71.6	3,480	3,496	75.9
Black/African American	124,032	123,789	1,244.4	135,270	134,886	1,312.4	147,238	146,651	1,383.7	159,674	158,656	1,453.8
Hispanic/Latino ^c	51,055	51,082	597.5	56,572	56,572	626.7	62,570	62,534	657.2	68,842	68,715	685.9
Native Hawaiian/Other Pacific Islander	207	208	201.5	228	229	209.7	256	257	222.8	285	286	235.2
White	114,101	113,797	150.4	126,146	125,717	163.2	138,578	137,979	176.2	151,928	151,037	190.1
Multiple races	9,846	9,844	1,241.1	10,770	10,751	1,295.8	11,625	11,576	1,336.1	12,560	12,462	1,379.6
Total^f	302,487	301,970	303.1	332,578	331,753	325.1	364,309	363,046	347.8	397,793	395,668	371.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 6b. Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate									
Males												
Age at end of year (yr)												
50–54	106,326	106,312	959.1	113,375	113,335	1,012.1	120,549	120,481	1,074.7	127,973	127,806	1,140.4
55–59	68,984	68,835	709.6	76,035	75,821	763.3	83,270	82,973	814.3	90,277	89,822	863.5
60–64	37,233	37,094	448.7	42,432	42,217	487.8	47,592	47,272	546.8	53,260	52,775	601.0
65–69	16,289	16,200	270.8	18,695	18,559	301.0	21,836	21,632	322.5	25,464	25,121	358.4
70–74	7,029	6,979	160.8	8,140	8,063	180.3	9,483	9,361	200.5	11,015	10,823	218.3
75–79	2,803	2,771	85.7	3,247	3,200	97.5	3,687	3,619	108.2	4,324	4,221	122.5
80–84	992	975	41.7	1,151	1,127	47.6	1,387	1,350	56.5	1,609	1,551	64.6
≥85	403	393	21.4	487	473	24.8	569	548	27.5	690	655	31.7
Race/ethnicity												
American Indian/Alaska Native	538	535	—	595	591	—	662	657	—	738	731	—
Asian ^b	2,041	2,047	—	2,249	2,256	—	2,574	2,583	—	2,872	2,885	—
Black/African American	84,477	84,243	—	91,650	91,300	—	99,084	98,567	—	106,886	106,043	—
Hispanic/Latino ^c	43,828	43,836	—	48,501	48,477	—	53,660	53,598	—	59,106	58,954	—
Native Hawaiian/Other Pacific Islander	169	170	—	188	189	—	211	212	—	230	231	—
White	101,577	101,304	—	112,277	111,897	—	123,434	122,912	—	135,273	134,503	—
Multiple races	7,429	7,424	—	8,102	8,084	—	8,748	8,707	—	9,507	9,428	—
Transmission category												
Male-to-male sexual contact	122,627	140,068	—	136,908	156,494	—	152,207	174,125	—	168,479	192,732	—
Injection drug use	41,961	49,880	—	43,615	52,026	—	45,201	54,092	—	46,795	56,053	—
Male-to-male sexual contact and injection drug use	16,827	18,457	—	18,454	20,218	—	20,112	22,015	—	21,890	23,893	—
Heterosexual contact ^d	23,085	30,065	—	25,298	32,898	—	27,520	35,771	—	29,913	38,800	—
Other ^e	35,559	1,089	—	39,287	1,159	—	43,333	1,233	—	47,535	1,296	—
Subtotal, males	240,059	239,559	512.1	263,562	262,794	547.7	288,373	287,236	584.5	314,612	312,775	622.2
Females												
Age at end of year (yr)												
50–54	31,710	31,740	275.2	33,757	33,785	290.1	35,777	35,797	307.2	37,693	37,684	323.6
55–59	19,795	19,791	191.4	22,174	22,152	209.3	24,572	24,532	226.2	27,080	26,991	244.2
60–64	10,122	10,114	112.9	11,700	11,679	124.2	13,362	13,315	141.5	15,072	14,981	156.4
65–69	4,406	4,398	65.4	5,088	5,074	73.4	5,911	5,885	78.6	6,983	6,928	88.6
70–74	1,992	1,983	38.6	2,294	2,281	43.1	2,718	2,698	49.0	3,138	3,101	53.3
75–79	880	875	20.9	1,011	1,001	23.7	1,162	1,147	26.9	1,361	1,338	30.7
80–84	339	335	9.6	387	382	10.9	478	469	13.5	550	537	15.6
≥85	162	158	4.2	180	174	4.5	195	185	4.7	230	216	5.3

Table 6b. Adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
Race/ethnicity												
American Indian/Alaska Native	214	213	—	242	241	—	261	260	—	287	286	—
Asian ^b	469	471	—	524	527	—	564	568	—	626	629	—
Black/African American	39,688	39,680	—	43,763	43,730	—	48,295	48,226	—	52,947	52,772	—
Hispanic/Latino ^c	14,020	14,044	—	15,445	15,462	—	16,946	16,952	—	18,437	18,421	—
Native Hawaiian/Other Pacific Islander	41	41	—	43	43	—	48	48	—	59	60	—
White	12,553	12,521	—	13,901	13,852	—	15,179	15,101	—	16,694	16,573	—
Multiple races	2,421	2,423	—	2,673	2,672	—	2,882	2,874	—	3,057	3,037	—
Transmission category												
Injection drug use	17,500	22,391	—	18,814	24,156	—	20,101	25,906	—	21,478	27,713	—
Heterosexual contact ^d	34,452	46,149	—	38,242	51,496	—	42,217	57,217	—	46,313	63,128	—
Other ^e	17,454	855	—	19,535	876	—	21,857	906	—	24,316	935	—
Subtotal, females	69,406	69,394	128.1	76,591	76,528	138.1	84,175	84,030	148.5	92,107	91,776	158.9
Both sexes												
Age at end of year (yr)												
50–54	138,036	138,051	610.3	147,132	147,120	644.0	156,326	156,278	683.6	165,666	165,490	724.1
55–59	88,779	88,626	442.2	98,209	97,973	477.6	107,842	107,504	511.0	117,357	116,813	544.4
60–64	47,355	47,208	274.1	54,132	53,896	298.5	60,954	60,587	335.5	68,332	67,757	369.0
65–69	20,695	20,599	162.1	23,783	23,633	180.6	27,747	27,518	193.8	32,447	32,050	216.2
70–74	9,021	8,961	94.5	10,434	10,344	106.0	12,201	12,060	118.6	14,153	13,924	129.2
75–79	3,683	3,646	49.1	4,258	4,201	56.0	4,849	4,766	62.6	5,685	5,559	71.2
80–84	1,331	1,310	22.5	1,538	1,508	25.8	1,865	1,819	31.1	2,159	2,087	35.7
≥85	565	551	9.8	667	647	11.2	764	733	12.3	920	871	14.3
Race/ethnicity												
American Indian/Alaska Native	752	749	—	837	832	—	923	917	—	1,025	1,016	—
Asian ^b	2,510	2,518	—	2,773	2,783	—	3,138	3,151	—	3,498	3,514	—
Black/African American	124,165	123,923	—	135,413	135,030	—	147,379	146,793	—	159,833	158,815	—
Hispanic/Latino ^c	57,848	57,879	—	63,946	63,939	—	70,606	70,550	—	77,543	77,374	—
Native Hawaiian/Other Pacific Islander	210	211	—	231	232	—	259	260	—	289	290	—
White	114,130	113,826	—	126,178	125,749	—	138,613	138,013	—	151,967	151,076	—
Multiple races	9,850	9,848	—	10,775	10,756	—	11,630	11,581	—	12,564	12,465	—
Total^f	309,465	308,953	306.1	340,153	339,322	328.2	372,548	371,265	351.1	406,719	404,551	374.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 7a. Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
50–64 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	492	490	260.4	539	536	276.1	592	588	297.1	656	650	321.8
Asian ^b	1,776	1,781	153.7	1,940	1,947	161.0	2,194	2,204	175.8	2,427	2,441	187.0
Black/African American	75,285	75,141	2,497.2	81,173	80,957	2,599.9	86,935	86,622	2,717.2	92,772	92,260	2,832.7
Hispanic/Latino ^c	34,667	34,685	1,245.9	38,328	38,332	1,301.8	42,210	42,197	1,362.1	46,316	46,258	1,420.6
Native Hawaiian/Other Pacific Islander	146	147	411.7	166	167	446.3	186	187	478.7	201	201	496.5
White	89,383	89,201	419.3	98,021	97,773	451.1	106,748	106,421	491.5	115,641	115,173	530.7
Multiple races	6,721	6,721	2,627.7	7,294	7,285	2,725.7	7,824	7,797	2,829.2	8,438	8,385	2,946.8
Transmission category												
Male-to-male sexual contact	107,812	123,081	—	119,701	136,819	—	132,074	151,170	—	145,060	166,090	—
Injection drug use	36,847	43,182	—	37,796	44,491	—	38,414	45,434	—	38,829	46,076	—
Male-to-male sexual contact and injection drug use	15,569	17,016	—	16,973	18,535	—	18,366	20,045	—	19,766	21,524	—
Heterosexual contact ^d	18,582	24,080	—	20,299	26,279	—	21,969	28,429	—	23,803	30,698	—
Other ^e	29,660	808	—	32,692	873	—	35,866	938	—	38,993	979	—
Subtotal, males	208,470	208,167	725.1	227,461	226,997	771.0	246,689	246,015	828.2	266,451	265,367	883.1
Females												
Race/ethnicity												
American Indian/Alaska Native	196	196	95.1	222	222	103.9	235	234	107.3	257	256	114.9
Asian ^b	383	385	27.7	428	431	29.7	464	468	31.1	508	511	32.8
Black/African American	35,384	35,396	1,003.5	38,847	38,845	1,064.1	42,550	42,529	1,138.2	46,167	46,079	1,207.6
Hispanic/Latino ^c	10,352	10,370	351.8	11,319	11,334	364.9	12,319	12,329	378.8	13,251	13,250	388.8
Native Hawaiian/Other Pacific Islander	38	38	104.3	38	38	99.7	42	42	105.3	53	54	127.9
White	11,216	11,194	50.8	12,337	12,303	54.7	13,320	13,266	59.1	14,522	14,434	64.1
Multiple races	2,205	2,208	790.4	2,424	2,425	829.8	2,601	2,596	859.6	2,736	2,723	871.9
Transmission category												
Injection drug use	16,228	20,377	—	17,330	21,842	—	18,330	23,190	—	19,348	24,521	—
Heterosexual contact ^d	28,698	38,827	—	31,704	43,163	—	34,771	47,667	—	37,823	52,168	—
Other ^e	14,848	583	—	16,581	592	—	18,430	606	—	20,323	617	—
Subtotal, females	59,774	59,787	196.5	65,615	65,598	210.1	71,531	71,463	226.8	77,494	77,306	242.5
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	688	685	174.1	761	758	185.9	827	822	197.6	913	906	213.3
Asian ^b	2,159	2,167	84.9	2,368	2,378	89.4	2,658	2,672	97.0	2,935	2,952	103.0
Black/African American	110,669	110,537	1,691.2	120,020	119,802	1,771.1	129,485	129,151	1,865.1	138,939	138,339	1,956.0
Hispanic/Latino ^c	45,019	45,055	786.1	49,647	49,665	820.8	54,529	54,525	858.3	59,567	59,508	893.0
Native Hawaiian/Other Pacific Islander	184	185	255.8	204	205	270.5	228	229	289.2	254	255	309.4
White	100,599	100,395	231.8	110,358	110,077	249.3	120,068	119,686	271.3	130,163	129,607	293.0
Multiple races	8,926	8,929	1,668.7	9,718	9,710	1,735.5	10,425	10,393	1,799.4	11,174	11,108	1,861.1
Subtotal, 50–64 years of age at end of year	268,244	267,954	453.1	293,076	292,594	482.3	318,220	317,478	518.6	343,945	342,674	553.3

Table 7a. Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
≥65 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	45	44	54.5	55	54	63.1	69	68	73.7	81	80	81.3
Asian ^b	252	252	41.3	294	294	45.1	364	363	51.9	429	428	56.8
Black/African American	9,106	9,015	672.6	10,382	10,247	739.5	12,058	11,854	813.3	14,013	13,682	893.9
Hispanic/Latino ^c	4,474	4,465	372.7	5,127	5,110	402.9	5,960	5,932	438.5	6,851	6,795	471.7
Native Hawaiian/Other Pacific Islander	20	20	143.3	19	19	125.1	22	22	132.7	25	25	138.9
White	12,172	12,081	85.6	14,231	14,099	97.8	16,658	16,463	109.1	19,603	19,302	123.5
Multiple races	706	701	627.8	805	796	676.6	921	907	716.9	1,067	1,042	772.8
Transmission category												
Male-to-male sexual contact	13,642	15,789	—	15,904	18,341	—	18,657	21,445	—	21,804	24,993	—
Injection drug use	3,183	4,700	—	3,787	5,433	—	4,592	6,397	—	5,599	7,546	—
Male-to-male sexual contact and injection drug use	920	1,103	—	1,112	1,313	—	1,343	1,567	—	1,683	1,928	—
Heterosexual contact ^d	3,277	4,717	—	3,683	5,258	—	4,170	5,917	—	4,635	6,581	—
Other ^e	5,753	271	—	6,427	275	—	7,290	285	—	8,348	306	—
Subtotal, males	26,775	26,579	152.1	30,913	30,619	170.7	36,052	35,610	189.1	42,069	41,354	210.9
Females												
Race/ethnicity												
American Indian/Alaska Native	18	18	17.5	20	20	18.4	26	25	22.4	30	30	24.6
Asian ^b	84	84	10.5	94	94	11.0	98	98	10.7	116	116	11.8
Black/African American	4,257	4,236	204.5	4,868	4,837	227.3	5,695	5,647	254.7	6,722	6,635	287.2
Hispanic/Latino ^c	1,562	1,562	96.5	1,798	1,796	105.2	2,081	2,077	114.7	2,424	2,412	126.0
Native Hawaiian/Other Pacific Islander	3	3	17.9	5	5	27.7	6	6	30.9	6	6	28.6
White	1,330	1,321	7.2	1,557	1,542	8.3	1,852	1,829	9.6	2,162	2,129	10.9
Multiple races	214	214	145.9	247	246	160.9	279	276	170.1	319	312	181.8
Transmission category												
Injection drug use	799	1522	—	959	1,770	—	1,199	2,125	—	1,494	2,536	—
Heterosexual contact ^d	4,144	5,653	—	4,766	6,495	—	5,507	7,541	—	6,406	8,794	—
Other ^e	2,525	262	—	2,864	274	—	3,331	292	—	3,879	310	—
Subtotal, females	7,468	7,437	32.3	8,589	8,540	36.4	10,037	9,958	40.9	11,779	11,639	46.3
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	63	62	34.0	75	74	38.3	95	93	45.4	111	109	50.0
Asian ^b	336	336	23.8	388	388	25.8	462	461	28.5	545	544	31.3
Black/African American	13,363	13,252	388.4	15,250	15,084	429.3	17,753	17,501	476.3	20,735	20,317	529.0
Hispanic/Latino ^c	6,036	6,027	213.9	6,925	6,907	232.1	8,041	8,009	253.2	9,275	9,207	274.5
Native Hawaiian/Other Pacific Islander	23	23	74.8	24	24	72.3	28	28	77.9	31	31	79.6
White	13,502	13,402	41.4	15,788	15,641	47.6	18,510	18,292	53.5	21,765	21,430	60.8
Multiple races	920	914	354.4	1,052	1,041	385.3	1,200	1,183	409.6	1,386	1,354	441.8
Subtotal, ≥65 years of age at end of year	34,243	34,017	84.0	39,502	39,159	94.7	46,089	45,568	105.6	53,848	52,994	118.5
Total^f	302,487	301,970	303.1	332,578	331,753	325.1	364,309	363,046	347.8	397,793	395,668	371.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 7b. Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
50–64 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	493	491	—	540	537	—	593	589	—	657	651	—
Asian ^b	1,785	1,790	—	1,951	1,958	—	2,206	2,216	—	2,439	2,453	—
Black/African American	75,353	75,210	—	81,249	81,033	—	87,004	86,691	—	92,847	92,335	—
Hispanic/Latino ^c	38,639	38,657	—	42,595	42,593	—	46,823	46,797	—	51,259	51,178	—
Native Hawaiian/Other Pacific Islander	149	150	—	169	170	—	189	190	—	205	205	—
White	89,401	89,219	—	98,041	97,793	—	106,769	106,442	—	115,663	115,195	—
Multiple races	6,723	6,724	—	7,297	7,288	—	7,827	7,800	—	8,440	8,386	—
Transmission category												
Male-to-male sexual contact	108,817	124,108	—	120,812	137,958	—	133,335	152,462	—	146,428	167,492	—
Injection drug use	38,611	44,990	—	39,645	46,388	—	40,386	47,452	—	40,921	48,213	—
Male-to-male sexual contact and injection drug use	15,882	17,330	—	17,314	18,878	—	18,737	20,417	—	20,171	21,930	—
Heterosexual contact ^d	19,472	25,001	—	21,256	27,271	—	22,956	29,452	—	24,851	31,784	—
Other ^e	29,761	811	—	32,815	877	—	35,997	942	—	39,139	983	—
Subtotal, males	212,543	212,241	730.6	231,842	231,372	776.8	251,411	250,725	834.5	271,510	270,403	889.8
Females												
Race/ethnicity												
American Indian/Alaska Native	196	196	—	222	222	—	235	234	—	257	256	—
Asian ^b	385	387	—	430	433	—	466	470	—	510	513	—
Black/African American	35,421	35,433	—	38,883	38,881	—	42,586	42,565	—	46,210	46,122	—
Hispanic/Latino ^c	12,157	12,180	—	13,288	13,306	—	14,453	14,464	—	15,546	15,544	—
Native Hawaiian/Other Pacific Islander	38	38	—	38	38	—	42	42	—	53	54	—
White	11,223	11,201	—	12,344	12,310	—	13,326	13,272	—	14,531	14,443	—
Multiple races	2,207	2,210	—	2,426	2,427	—	2,603	2,598	—	2,738	2,725	—
Transmission category												
Injection drug use	16,669	20,833	—	17,818	22,346	—	18,860	23,736	—	19,934	25,121	—
Heterosexual contact ^d	30,048	40,224	—	33,162	44,675	—	36,347	49,298	—	39,506	53,915	—
Other ^e	14,910	588	—	16,651	597	—	18,504	610	—	20,405	620	—
Subtotal, females	61,627	61,645	199.9	67,631	67,617	213.8	73,711	73,645	230.8	79,845	79,656	246.8
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	689	686	—	762	759	—	828	823	—	914	907	—
Asian ^b	2,170	2,178	—	2,381	2,391	—	2,672	2,686	—	2,949	2,966	—
Black/African American	110,774	110,643	—	120,132	119,914	—	129,590	129,256	—	139,057	138,457	—
Hispanic/Latino ^c	50,796	50,837	—	55,883	55,898	—	61,276	61,261	—	66,805	66,722	—
Native Hawaiian/Other Pacific Islander	187	188	—	207	208	—	231	232	—	258	259	—
White	100,624	100,420	—	110,385	110,104	—	120,095	119,713	—	130,194	129,638	—
Multiple races	8,930	8,933	—	9,723	9,715	—	10,430	10,398	—	11,178	11,111	—
Subtotal, 50–64 years of age at end of year	274,170	273,885	457.4	299,473	298,989	486.8	325,122	324,370	523.6	351,355	350,060	558.6

Table 7b. Adults aged 50 years and older living with diagnosed HIV infection, by year, age group, and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
≥65 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	45	44	—	55	54	—	69	68	—	81	80	—
Asian ^b	256	256	—	298	298	—	368	367	—	433	432	—
Black/African American	9,124	9,033	—	10,401	10,266	—	12,080	11,876	—	14,039	13,708	—
Hispanic/Latino ^c	5,189	5,178	—	5,906	5,885	—	6,837	6,800	—	7,847	7,776	—
Native Hawaiian/Other Pacific Islander	20	20	—	19	19	—	22	22	—	25	25	—
White	12,176	12,085	—	14,236	14,104	—	16,665	16,470	—	19,610	19,308	—
Multiple races	706	701	—	805	796	—	921	907	—	1,067	1,042	—
Transmission category												
Male-to-male sexual contact	13,810	15,960	—	16,096	18,536	—	18,872	21,663	—	22,051	25,240	—
Injection drug use	3,350	4,890	—	3,970	5,637	—	4,815	6,640	—	5,874	7,840	—
Male-to-male sexual contact and injection drug use	945	1,127	—	1,140	1,341	—	1,375	1,598	—	1,719	1,963	—
Heterosexual contact ^d	3,613	5,064	—	4,042	5,626	—	4,564	6,319	—	5,062	7,016	—
Other ^e	5,798	278	—	6,472	282	—	7,336	290	—	8,396	312	—
Subtotal, males	27,516	27,318	154.1	31,720	31,421	172.7	36,962	36,510	191.2	43,102	42,371	213.1
Females												
Race/ethnicity												
American Indian/Alaska Native	18	18	—	20	20	—	26	25	—	30	30	—
Asian ^b	84	84	—	94	94	—	98	98	—	116	116	—
Black/African American	4,267	4,246	—	4,880	4,849	—	5,709	5,661	—	6,737	6,650	—
Hispanic/Latino ^c	1,863	1,864	—	2,157	2,156	—	2,493	2,489	—	2,891	2,877	—
Native Hawaiian/Other Pacific Islander	3	3	—	5	5	—	6	6	—	6	6	—
White	1,330	1,321	—	1,557	1,542	—	1,853	1,830	—	2,163	2,130	—
Multiple races	214	214	—	247	246	—	279	276	—	319	312	—
Transmission category												
Injection drug use	831	1,558	—	996	1,810	—	1,241	2,171	—	1,544	2,592	—
Heterosexual contact ^d	4,404	5,925	—	5,080	6,821	—	5,870	7,919	—	6,807	9,213	—
Other ^e	2,544	267	—	2,884	280	—	3,353	296	—	3,911	315	—
Subtotal, females	7,779	7,749	33.2	8,960	8,911	37.5	10,464	10,385	42.1	12,262	12,120	47.6
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	63	62	—	75	74	—	95	93	—	111	109	—
Asian ^b	340	340	—	392	392	—	466	465	—	549	548	—
Black/African American	13,391	13,280	—	15,281	15,115	—	17,789	17,537	—	20,776	20,358	—
Hispanic/Latino ^c	7,052	7,042	—	8,063	8,041	—	9,330	9,289	—	10,738	10,653	—
Native Hawaiian/Other Pacific Islander	23	23	—	24	24	—	28	28	—	31	31	—
White	13,506	13,405	—	15,793	15,645	—	18,518	18,300	—	21,773	21,438	—
Multiple races	920	914	—	1,052	1,041	—	1,200	1,183	—	1,386	1,354	—
Subtotal, ≥65 years of age at end of year	35,295	35,068	85.4	40,680	40,333	96.1	47,426	46,896	107.1	55,364	54,492	120.1
Total^f	309,465	308,953	306.1	340,153	339,322	328.2	372,548	371,265	351.1	406,719	404,551	374.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 8a. Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States

	Age at end of year											
	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	349	346	441.3	202	200	290.1	105	104	189.9	46	45	113.1
Asian ^b	1,139	1,148	230.6	799	802	181.4	489	491	134.3	263	263	96.0
Black/African American	41,596	41,537	3,260.0	32,111	31,900	2,841.0	19,065	18,824	2,188.9	8,327	8,161	1,378.0
Hispanic/Latino ^c	23,295	23,312	1,657.7	14,974	14,945	1,389.0	8,047	8,001	1,033.8	3,863	3,839	704.4
Native Hawaiian/Other Pacific Islander	85	85	514.1	75	76	558.6	41	41	389.0	11	11	145.1
White	54,990	54,877	712.8	37,665	37,481	502.0	22,986	22,815	348.9	11,762	11,629	215.3
Multiple races	4,156	4,142	3,655.5	2,737	2,714	2,782.8	1,545	1,528	2,073.8	676	664	1,251.4
Transmission category												
Male-to-male sexual contact	72,526	82,936	—	46,029	52,898	—	26,505	30,256	—	13,120	15,005	—
Injection drug use	14,196	17,322	—	14,710	17,273	—	9,923	11,481	—	3,822	4,763	—
Male-to-male sexual contact and injection drug use	9,986	10,884	—	6,579	7,150	—	3,201	3,490	—	1,180	1,320	—
Heterosexual contact ^d	10,971	13,836	—	8,093	10,500	—	4,739	6,362	—	2,449	3,413	—
Other ^e	17,931	469	—	13,152	297	—	7,910	214	—	4,377	113	—
Subtotal, males	125,610	125,447	1,131.7	88,563	88,117	856.5	52,278	51,803	597.0	24,948	24,613	356.1
Females												
Race/ethnicity												
American Indian/Alaska Native	120	120	141.0	82	81	105.3	55	55	90.6	18	18	40.0
Asian ^b	206	207	36.1	186	187	35.3	116	117	25.7	54	54	16.1
Black/African American	21,584	21,594	1,494.1	15,823	15,778	1,203.5	8,760	8,706	821.9	3,828	3,793	496.5
Hispanic/Latino ^c	6,352	6,364	450.8	4,384	4,382	385.8	2,515	2,505	291.0	1,375	1,371	213.7
Native Hawaiian/Other Pacific Islander	21	21	126.6	20	20	143.5	12	12	110.2	5	5	63.5
White	7,044	7,014	89.4	4,864	4,829	62.3	2,614	2,591	37.4	1,262	1,248	21.4
Multiple races	1,270	1,267	1,019.6	917	911	856.8	549	545	667.3	176	173	285.3
Transmission category												
Injection drug use	8,913	11,331	—	6,907	8,654	—	3,528	4,537	—	1,099	1,641	—
Heterosexual contact ^d	18,162	25,026	—	12,574	17,334	—	7,087	9,809	—	3,579	4,907	—
Other ^e	9,522	230	—	6,795	201	—	4,006	185	—	2,040	115	—
Subtotal, females	36,597	36,587	318.0	26,276	26,188	239.8	14,621	14,531	153.7	6,718	6,663	86.5
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	469	466	285.0	284	281	192.6	160	159	137.6	64	63	74.6
Asian ^b	1,345	1,355	126.4	985	989	101.7	605	607	74.1	317	317	51.9
Black/African American	63,180	63,131	2,321.5	47,934	47,678	1,958.9	27,825	27,530	1,434.5	12,155	11,955	881.4
Hispanic/Latino ^c	29,647	29,676	1,053.1	19,358	19,327	873.8	10,562	10,505	642.7	5,238	5,210	439.1
Native Hawaiian/Other Pacific Islander	106	106	319.2	95	96	347.2	53	53	246.1	16	16	103.6
White	62,034	61,891	398.1	42,529	42,310	278.1	25,600	25,406	188.6	13,024	12,877	114.5
Multiple races	5,426	5,409	2,277.0	3,654	3,625	1,778.2	2,094	2,073	1,334.2	852	837	736.6
Total^f	162,207	162,034	717.2	114,839	114,306	538.9	66,899	66,334	365.8	31,666	31,276	214.0

Table 8a. Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States (cont)

	Age at end of year											
	70–74 years			75–79 years			80–84 years			≥85 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	23	23	86.6	8	8	48.5	2	2	21.0	2	2	30.3
Asian ^b	98	98	50.3	50	50	37.3	13	13	15.3	5	5	7.0
Black/African American	3,523	3,440	871.7	1,374	1,329	507.5	535	512	325.4	254	240	192.8
Hispanic/Latino ^c	1,806	1,793	500.5	747	738	298.2	311	305	188.8	124	121	94.0
Native Hawaiian/Other Pacific Islander	8	8	171.9	3	3	102.9	2	2	112.1	1	1	93.1
White	5,020	4,935	127.4	1,901	1,859	68.6	661	635	32.7	259	244	14.4
Multiple races	247	240	693.0	103	99	441.3	29	28	200.6	12	11	103.2
Transmission category												
Male-to-male sexual contact	5,563	6,393	—	2,111	2,448	—	746	848	—	264	300	—
Injection drug use	1,248	1,809	—	353	612	—	118	248	—	58	113	—
Male-to-male sexual contact and injection drug use	367	429	—	92	120	—	37	49	—	7	11	—
Heterosexual contact ^d	1,290	1,826	—	596	860	—	202	324	—	98	159	—
Other ^e	2,257	80	—	1,034	46	—	450	27	—	230	41	—
Subtotal, males	10,725	10,536	215.7	4,186	4,085	120.4	1,553	1,496	63.1	657	624	30.5
Females												
Race/ethnicity												
American Indian/Alaska Native	8	8	26.0	2	2	9.8	1	1	7.7	1	1	7.7
Asian ^b	38	38	16.1	10	10	5.8	8	8	6.5	6	6	4.9
Black/African American	1,702	1,678	305.1	784	770	190.2	298	291	101.5	110	103	33.8
Hispanic/Latino ^c	645	642	140.0	261	258	75.9	103	101	41.6	40	39	17.0
Native Hawaiian/Other Pacific Islander	0	0	0.1	0	0	0.0	0	0	0.0	1	1	48.2
White	530	523	11.9	214	209	6.3	99	96	3.5	57	53	1.6
Multiple races	86	84	203.4	38	37	130.9	17	16	80.4	2	1	6.0
Transmission category												
Injection drug use	292	577	—	64	195	—	28	86	—	11	37	—
Heterosexual contact ^d	1,673	2,317	—	774	1,041	—	286	398	—	94	130	—
Other ^e	1,044	78	—	471	50	—	212	29	—	112	37	—
Subtotal, females	3,009	2,973	51.9	1,309	1,286	30.0	526	514	15.1	217	204	5.1
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	31	31	54.3	10	10	26.9	3	3	13.1	3	3	15.6
Asian ^b	136	136	31.5	60	60	19.6	21	21	10.0	11	11	5.7
Black/African American	5,225	5,118	541.8	2,158	2,099	314.8	833	803	180.8	364	343	79.9
Hispanic/Latino ^c	2,451	2,435	298.1	1,008	996	169.6	414	406	100.4	164	160	44.6
Native Hawaiian/Other Pacific Islander	8	8	81.6	3	3	46.6	2	2	48.5	2	2	63.5
White	5,550	5,457	65.9	2,115	2,068	34.3	760	731	15.7	316	297	5.9
Multiple races	333	324	426.3	141	136	267.5	46	44	129.1	14	13	39.5
Total^f	13,734	13,509	127.3	5,495	5,372	69.9	2,079	2,009	34.8	874	828	13.7

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 8b. Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States and 6 dependent areas

	Age at end of year											
	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	350	347	—	202	200	—	105	104	—	46	45	—
Asian ^b	1,145	1,154	—	803	806	—	491	493	—	265	265	—
Black/African American	41,628	41,569	—	32,139	31,927	—	19,080	18,839	—	8,345	8,179	—
Hispanic/Latino ^c	25,605	25,618	—	16,649	16,610	—	9,005	8,949	—	4,356	4,325	—
Native Hawaiian/Other Pacific Islander	87	87	—	75	76	—	43	43	—	11	11	—
White	55,000	54,887	—	37,672	37,488	—	22,991	22,820	—	11,765	11,632	—
Multiple races	4,158	4,144	—	2,737	2,714	—	1,545	1,528	—	676	664	—
Transmission category												
Male-to-male sexual contact	73,230	83,661	—	46,453	53,332	—	26,745	30,499	—	13,246	15,132	—
Injection drug use	15,153	18,303	—	15,465	18,046	—	10,303	11,864	—	4,001	4,946	—
Male-to-male sexual contact and injection drug use	10,185	11,084	—	6,701	7,273	—	3,285	3,573	—	1,206	1,345	—
Heterosexual contact ^d	11,401	14,288	—	8,452	10,872	—	4,998	6,625	—	2,616	3,584	—
Other ^e	18,004	470	—	13,206	299	—	7,929	215	—	4,395	115	—
Subtotal, males	127,973	127,806	1,140.4	90,277	89,822	863.5	53,260	52,775	601.0	25,464	25,121	358.4
Females												
Race/ethnicity												
American Indian/Alaska Native	120	120	—	82	81	—	55	55	—	18	18	—
Asian ^b	207	208	—	186	187	—	117	118	—	54	54	—
Black/African American	21,603	21,613	—	15,837	15,793	—	8,770	8,716	—	3,836	3,802	—
Hispanic/Latino ^c	7,425	7,438	—	5,171	5,167	—	2,950	2,939	—	1,631	1,628	—
Native Hawaiian/Other Pacific Islander	21	21	—	20	20	—	12	12	—	5	5	—
White	7,047	7,017	—	4,866	4,831	—	2,618	2,595	—	1,263	1,249	—
Multiple races	1,270	1,267	—	918	912	—	550	546	—	176	173	—
Transmission category												
Injection drug use	9,218	11,642	—	7,093	8,845	—	3,623	4,633	—	1,129	1,675	—
Heterosexual contact ^d	18,916	25,811	—	13,159	17,943	—	7,431	10,160	—	3,796	5,136	—
Other ^e	9,559	230	—	6,828	202	—	4,018	188	—	2,058	117	—
Subtotal, females	37,693	37,684	323.6	27,080	26,991	244.2	15,072	14,981	156.4	6,983	6,928	88.6
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	470	467	—	284	281	—	160	159	—	64	63	—
Asian ^b	1,352	1,362	—	989	993	—	608	610	—	319	319	—
Black/African American	63,231	63,182	—	47,976	47,720	—	27,850	27,555	—	12,181	11,981	—
Hispanic/Latino ^c	33,030	33,056	—	21,820	21,778	—	11,955	11,888	—	5,987	5,952	—
Native Hawaiian/Other Pacific Islander	108	108	—	95	96	—	55	55	—	16	16	—
White	62,047	61,904	—	42,538	42,319	—	25,609	25,415	—	13,028	12,881	—
Multiple races	5,428	5,411	—	3,655	3,626	—	2,095	2,074	—	852	837	—
Total^f	165,666	165,490	724.1	117,357	116,813	544.4	68,332	67,757	369.0	32,447	32,050	216.2

Table 8b. Adults aged 50 years and older living with diagnosed HIV infection, by 5-year age group and selected characteristics, year-end 2013—United States and 6 dependent areas (cont)

	Age at end of year											
	70–74 years			75–79 years			80–84 years			≥85 years		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	23	23	—	8	8	—	2	2	—	2	2	—
Asian ^b	100	100	—	50	50	—	13	13	—	5	5	—
Black/African American	3,530	3,447	—	1,374	1,329	—	536	513	—	254	240	—
Hispanic/Latino ^c	2,086	2,070	—	883	872	—	365	357	—	157	152	—
Native Hawaiian/Other Pacific Islander	8	8	—	3	3	—	2	2	—	1	1	—
White	5,021	4,935	—	1,903	1,861	—	662	636	—	259	244	—
Multiple races	247	240	—	103	99	—	29	28	—	12	11	—
Transmission category												
Male-to-male sexual contact	5,635	6,465	—	2,146	2,482	—	753	855	—	271	306	—
Injection drug use	1,307	1,874	—	375	638	—	126	258	—	65	124	—
Male-to-male sexual contact and injection drug use	371	432	—	96	123	—	38	50	—	8	12	—
Heterosexual contact ^d	1,432	1,971	—	666	931	—	239	361	—	109	170	—
Other ^e	2,270	81	—	1,041	47	—	453	27	—	237	43	—
Subtotal, males	11,015	10,823	218.3	4,324	4,221	122.5	1,609	1,551	64.6	690	655	31.7
Females												
Race/ethnicity												
American Indian/Alaska Native	8	8	—	2	2	—	1	1	—	1	1	—
Asian ^b	38	38	—	10	10	—	8	8	—	6	6	—
Black/African American	1,705	1,681	—	786	772	—	299	292	—	111	104	—
Hispanic/Latino ^c	771	767	—	311	308	—	126	123	—	52	51	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	1	1	—
White	530	523	—	214	209	—	99	96	—	57	53	—
Multiple races	86	84	—	38	37	—	17	16	—	2	1	—
Transmission category												
Injection drug use	307	593	—	67	198	—	30	89	—	11	37	—
Heterosexual contact ^d	1,782	2,428	—	820	1,090	—	304	418	—	105	141	—
Other ^e	1,049	80	—	474	50	—	216	29	—	114	38	—
Subtotal, females	3,138	3,101	53.3	1,361	1,338	30.7	550	537	15.6	230	216	5.3
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	31	31	—	10	10	—	3	3	—	3	3	—
Asian ^b	138	138	—	60	60	—	21	21	—	11	11	—
Black/African American	5,235	5,128	—	2,160	2,100	—	835	805	—	365	344	—
Hispanic/Latino ^c	2,857	2,837	—	1,194	1,180	—	491	481	—	209	203	—
Native Hawaiian/Other Pacific Islander	8	8	—	3	3	—	2	2	—	2	2	—
White	5,551	5,458	—	2,117	2,070	—	761	732	—	316	297	—
Multiple races	333	324	—	141	136	—	46	44	—	14	13	—
Total^f	14,153	13,924	129.2	5,685	5,559	71.2	2,159	2,087	35.7	920	871	14.3

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information is limited for U.S. dependent areas. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 9. Adults aged 50 years and older living with diagnosed HIV infection, by age group and area of residence, year-end 2013—United States and 6 dependent areas

Area of residence	Age at year-end 2013								
	50–64 years			≥65 years			Total ≥50 years		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No. ^b	Rate
Alabama	3,778	3,692	381.2	471	442	61.2	4,249	4,134	244.6
Alaska	233	229	156.2	41	40	60.9	274	269	126.6
Arizona	4,755	4,732	391.6	685	674	66.0	5,440	5,406	242.6
Arkansas	1,662	1,641	287.8	217	212	46.6	1,879	1,853	180.8
California	46,498	46,488	658.4	7,917	7,831	163.0	54,415	54,319	457.8
Colorado	5,056	5,040	489.6	783	777	120.1	5,839	5,817	347.0
Connecticut	4,681	4,625	605.6	731	709	130.1	5,412	5,334	407.5
Delaware	1,264	1,254	665.8	177	174	117.9	1,441	1,428	425.1
District of Columbia	5,757	5,790	5,402.5	950	948	1,292.2	6,707	6,738	3,732.8
Florida	37,996	37,849	965.6	6,501	6,409	175.6	44,497	44,259	584.6
Georgia	12,615	12,785	689.7	1,691	1,691	141.4	14,306	14,476	474.7
Hawaii	1,118	1,116	408.9	221	219	99.5	1,339	1,335	270.9
Idaho	331	330	109.5	28	28	12.4	359	357	68.2
Illinois	11,962	11,814	468.9	1,809	1,739	99.7	13,771	13,553	317.9
Indiana	3,122	3,103	238.6	388	381	41.7	3,510	3,484	157.3
Iowa	668	667	107.2	106	105	21.8	774	771	70.0
Kansas	885	879	157.5	111	107	26.5	996	986	102.4
Kentucky	1,922	1,888	211.8	236	229	36.0	2,158	2,117	138.8
Louisiana	5,920	5,886	650.6	730	716	116.6	6,650	6,601	434.8
Maine	527	525	167.5	92	91	38.9	619	617	112.4
Maryland	11,801	12,008	996.2	1,752	1,782	224.3	13,553	13,791	689.5
Massachusetts	7,960	8,089	587.0	1,138	1,153	116.6	9,098	9,241	390.5
Michigan	5,269	5,205	248.9	737	710	47.7	6,006	5,914	165.3
Minnesota	2,564	2,555	231.4	341	336	44.5	2,905	2,892	155.4
Mississippi	2,643	2,584	449.7	361	343	82.4	3,004	2,927	295.4
Missouri	4,138	4,109	336.1	508	498	54.9	4,646	4,607	216.3
Montana	147	146	65.2	16	16	9.5	163	161	41.6
Nebraska	557	556	153.9	63	62	23.4	620	618	98.8
Nevada	2,579	2,561	485.1	437	428	112.4	3,016	2,989	328.8
New Hampshire	519	515	167.9	47	46	22.4	566	560	110.0
New Jersey	16,423	16,412	902.4	2,546	2,512	195.6	18,969	18,923	610.0
New Mexico	935	931	225.5	149	146	47.6	1,084	1,078	149.7
New York	55,043	54,947	1,409.0	10,710	10,621	375.1	65,753	65,568	974.1
North Carolina	9,154	9,019	471.5	1,182	1,136	80.7	10,336	10,155	305.9
North Dakota	67	66	47.5	10	10	9.6	77	76	31.4
Ohio	6,587	6,569	272.1	820	808	46.1	7,407	7,376	177.0
Oklahoma	1,751	1,724	235.2	218	211	38.3	1,969	1,934	150.9
Oregon	2,073	2,067	256.4	322	317	52.3	2,395	2,384	168.7
Pennsylvania	12,672	12,528	463.7	1,874	1,818	87.0	14,546	14,347	299.4
Rhode Island	873	864	392.6	112	110	67.5	985	973	254.3

Table 9. Adults aged 50 years and older living with diagnosed HIV infection, by age group and area of residence, year-end 2013—United States and 6 dependent areas (cont)

Area of residence	Age at year-end 2013								
	50–64 years			≥65 years			Total ≥50 years		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No. ^b	Rate
South Carolina	5,357	5,329	558.3	751	740	101.7	6,108	6,069	360.8
South Dakota	171	171	100.8	31	31	24.4	202	201	68.2
Tennessee	5,010	4,847	374.2	635	588	61.7	5,645	5,435	241.8
Texas	22,657	22,500	485.7	2,859	2,784	93.6	25,516	25,284	332.5
Utah	933	931	218.8	119	119	41.9	1,052	1,050	148.1
Vermont	208	206	141.1	30	28	27.7	238	234	94.4
Virginia	8,115	7,936	486.6	1,180	1,117	101.0	9,295	9,053	330.8
Washington	4,278	4,274	307.0	648	642	67.5	4,926	4,916	209.8
West Virginia	618	613	150.8	76	75	23.6	694	688	94.8
Wisconsin	1,997	1,985	164.8	278	274	32.2	2,275	2,259	109.9
Wyoming	96	96	78.1	13	13	16.3	109	109	53.9
Subtotal	343,945	342,674	553.3	53,848	52,994	118.5	397,793	395,668	371.0
U.S. dependent areas									
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0
Guam	25	25	101.5	5	5	37.7	30	30	79.2
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0
Puerto Rico	7,156	7,131	1,064.5	1,453	1,435	239.1	8,609	8,567	674.4
Republic of Palau	1	1	31.7	0	0	0.0	1	1	21.7
U.S. Virgin Islands	228	229	987.8	58	57	308.8	286	286	685.4
Subtotal	7,410	7,386	1,002.1	1,516	1,498	234.6	8,926	8,883	645.9
Total^d	351,355	350,060	558.6	55,364	54,492	120.1	406,719	404,551	374.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 10. Adults aged 50 years and older living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total ^c				
	Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d				
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.
Alabama	55	54	218.3	4	4	33.8	28	27	466.5	25	25	350.3	0	0	0.0	154	152	97.0	8	7	133.0	274	269	126.6		
Alaska	1	1	9.4	8	8	55.7	2,562	2,488	679.2	76	75	330.2	2	2	456.6	1,450	1,414	111.8	150	146	1,157.6	4,249	4,134	244.6		
Arizona	2	2	28.0	4	4	45.2	697	687	561.4	61	61	254.4	0	0	0.0	1,054	1,039	121.8	61	60	635.5	1,879	1,853	180.8		
Arkansas	151	151	245.2	31	31	62.0	493	488	761.5	1,079	1,075	302.4	8	8	325.7	3,599	3,574	213.0	79	79	468.6	5,440	5,406	242.6		
California	170	169	298.3	1,402	1,410	82.5	9,426	9,394	1,366.5	13,029	13,075	474.8	75	75	202.3	29,383	29,280	453.3	930	916	568.2	54,415	54,319	457.8		
Colorado	22	22	235.7	22	22	57.8	698	695	1,304.4	793	791	403.2	4	4	302.3	4,206	4,191	307.1	94	93	678.0	5,839	5,817	347.0		
Connecticut	5	5	217.8	25	25	75.5	1,807	1,780	1,857.0	1,509	1,489	1,630.8	0	0	0.0	1,965	1,936	179.5	101	100	1,317.7	5,412	5,334	407.5		
Delaware	6	6	1,215.3	27	27	649.8	4,924	4,944	4,594.5	317	319	2,886.6	5	5	9,403.3	1,244	1,254	2,279.5	184	183	8,699.5	6,707	6,738	3,732.8		
District of Columbia	1	1	78.6	1	1	12.7	910	900	1,616.5	79	79	744.1	0	0	0.0	415	412	159.7	35	35	1,495.3	1,441	1,428	425.1		
Florida	26	26	134.7	97	97	63.8	20,534	20,426	2,537.1	8,482	8,453	682.7	13	13	416.4	14,459	14,369	271.0	886	875	1,766.4	44,497	44,259	584.6		
Georgia	4	4	51.8	45	46	55.0	9,396	9,515	1,219.2	544	552	518.4	4	4	368.0	3,745	3,785	184.8	568	570	2,657.2	14,306	14,476	474.7		
Hawaii	4	4	443.0	202	203	83.2	54	53	1,261.3	93	93	437.6	106	106	308.2	812	808	607.4	68	67	121.4	1,339	1,335	270.9		
Idaho	0	0	0.0	6	6	54.3	110	110	628.4	41	41	204.4	0	0	0.1	605	603	57.6	12	12	290.5	774	771	70.0		
Illinois	3	3	58.9	2	2	38.4	13	13	823.2	36	36	140.5	0	0	0.0	298	296	61.5	7	7	151.3	359	357	68.2		
Indiana	18	18	285.5	116	116	66.7	6,179	6,034	1,134.7	1,922	1,900	552.1	5	5	605.4	4,868	4,824	151.5	663	656	2,825.4	13,771	13,553	317.9		
Iowa	1	1	19.3	19	19	84.7	1,044	1,036	666.1	196	196	350.7	0	0	0.0	2,154	2,138	108.9	96	94	768.4	3,510	3,484	157.3		
Kansas	3	3	45.4	7	7	42.2	218	216	503.1	107	106	248.2	0	0	0.0	623	616	72.8	38	37	445.4	996	986	102.4		
Kentucky	2	2	58.0	6	6	53.9	639	624	658.4	74	73	425.2	1	1	275.8	1,376	1,351	97.3	60	59	635.5	2,158	2,117	138.8		
Louisiana	8	8	95.7	20	20	102.2	3,954	3,924	970.3	276	275	659.9	0	0	0.0	2,295	2,279	220.4	97	95	973.1	6,650	6,601	434.8		
Maine	8	8	204.6	81	83	93.5	2,602	2,656	2,283.0	1,908	1,938	1,660.5	1	1	159.1	4,363	4,418	218.1	135	137	968.6	9,098	9,241	390.5		
Maryland	11	11	224.9	54	56	54.9	10,151	10,338	2,011.6	467	475	603.5	3	3	399.9	2,292	2,323	181.1	575	583	3,390.3	13,553	13,791	689.5		
Massachusetts	3	3	111.1	0	0	0.0	54	54	2,065.9	25	25	740.0	0	0	0.0	533	531	99.7	4	4	100.4	619	617	112.4		
Michigan	13	13	75.8	12	12	20.1	3,112	3,055	766.5	244	241	326.1	0	0	0.0	2,468	2,439	81.3	157	155	572.5	6,006	5,914	165.3		
Minnesota	30	29	217.9	28	28	71.7	796	791	1,579.6	190	190	625.1	1	1	224.6	1,818	1,810	105.4	42	42	427.1	2,905	2,892	155.4		
Mississippi	5	5	58.8	11	11	47.1	1,594	1,582	828.4	213	212	623.5	1	1	98.9	2,714	2,691	145.0	108	105	616.0	4,646	4,607	216.3		
Missouri	2	2	54.5	7	7	99.2	2,008	1,951	656.9	55	53	447.3	1	1	444.9	812	795	119.3	119	118	2,417.0	3,004	2,927	295.4		
Montana	9	9	63.7	0	0	0.0	2	2	255.5	3	3	52.8	0	0	0.0	145	144	39.8	4	4	85.9	163	161	41.6		
Nebraska	52	50	149.4	28	28	55.2	6,634	6,498	1,049.6	344	343	369.7	2	2	184.6	3,032	2,993	119.6	244	241	1,184.1	10,336	10,155	305.9		
Nevada	3	3	42.0	0	0	0.0	12	12	929.6	5	5	247.8	0	0	0.0	54	53	23.2	3	3	236.3	77	76	31.4		
New Hampshire	5	5	155.0	2	2	28.8	132	132	694.2	63	63	279.1	1	1	489.9	410	409	71.6	7	7	247.7	620	618	98.8		
New Jersey	0	0	0.0	5	5	76.8	55	55	1,516.0	64	63	940.9	0	0	0.0	429	424	86.8	13	13	422.6	566	560	110.0		
New Mexico	6	6	134.6	125	126	58.3	9,096	9,057	2,676.3	4,125	4,125	1,179.9	3	3	439.6	4,488	4,484	206.2	1,126	1,123	5,699.1	18,969	18,923	610.0		
New York	50	50	113.1	0	0	0.0	47	47	408.7	430	429	169.2	0	0	0.0	536	532	134.1	21	21	365.4	1,084	1,078	149.7		
North Carolina	19	19	247.5	51	51	68.2	630	622	1,006.1	468	467	378.0	4	4	106.1	1,786	1,769	282.7	58	57	506.8	3,016	2,989	328.8		
North Dakota	36	36	205.3	574	576	125.8	26,331	26,254	3,007.8	20,846	20,784	2,564.2	6	6	297.9	14,500	14,468	320.0	3,460	3,445	6,985.2	65,753	65,568	974.1		

Table 10. Adults aged 50 years and older living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2013—United States and 6 dependent areas (cont)

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total ^c					
	Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d			Estimated ^d					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Ohio	7	7	83.6	16	16	32.9	2,973	2,958	724.2	363	362	597.2	0	0	0.0	3,848	3,833	106.1	200	200	782.6	7,407	7,376	177.0			
Oklahoma	102	99	124.3	10	10	58.9	382	375	506.3	123	122	268.5	0	0	0.0	1,281	1,259	122.5	71	69	184.1	1,969	1,934	150.9			
Oregon	14	14	106.1	17	17	40.6	151	151	864.3	182	182	295.1	5	5	206.6	1,994	1,984	157.8	32	31	163.5	2,395	2,384	168.7			
Pennsylvania	7	7	112.2	52	52	56.3	6,722	6,617	1,719.7	2,089	2,064	1,620.9	9	9	1,200.7	5,097	5,036	121.1	570	561	2,545.0	14,546	14,347	299.4			
Rhode Island	5	5	371.8	7	7	99.7	227	223	1,559.5	219	218	949.0	1	1	716.9	508	502	150.2	18	17	660.2	985	973	254.3			
South Carolina	6	6	104.4	8	8	46.3	4,213	4,186	1,066.4	143	143	454.0	1	1	211.3	1,632	1,622	132.3	105	104	1,126.9	6,108	6,069	360.8			
South Dakota	30	29	225.6	2	2	113.8	28	28	1,009.0	6	6	157.1	0	0	0.0	134	134	49.3	2	2	103.9	202	201	68.2			
Tennessee	3	3	44.1	13	12	52.4	2,894	2,771	947.4	143	142	399.3	4	4	654.4	2,465	2,383	127.1	123	120	790.5	5,645	5,435	241.8			
Texas	22	22	74.8	132	132	48.5	8,254	8,171	1,028.2	6,132	6,103	319.5	6	6	147.4	10,188	10,079	221.9	782	771	1,501.0	25,516	25,284	332.5			
Utah	11	11	193.9	5	5	35.6	74	74	1,484.4	137	137	275.8	0	0	0.0	812	811	129.3	13	13	320.0	1,052	1,050	148.1			
Vermont	7	7	93.1	66	65	52.0	5,223	5,063	1,084.5	453	446	438.8	2	2	161.7	3,309	3,241	161.2	235	228	973.4	9,295	9,053	330.8			
Virginia	0	0	0.0	0	0	0.0	19	19	1,683.0	11	11	552.2	0	0	0.0	204	201	83.3	4	4	185.0	238	234	94.4			
Washington	59	59	230.5	107	107	71.0	656	656	1,045.3	418	419	402.1	11	11	137.7	3,565	3,554	181.6	110	110	327.7	4,926	4,916	209.8			
West Virginia	12	12	94.4	23	23	103.1	724	715	914.2	204	202	447.3	0	0	0.0	1,267	1,261	66.8	45	45	522.7	2,275	2,259	109.9			
Wisconsin	1	1	66.3	0	0	0.0	188	186	927.0	21	21	431.6	0	0	0.0	452	449	64.9	32	32	670.7	694	688	94.8			
Wyoming	4	4	148.6	0	0	0.0	4	4	129.2	9	9	82.2	0	0	0.0	87	87	47.6	5	5	334.5	109	109	53.9			
Subtotal	1,024	1,015	157.8	3,480	3,496	75.9	159,674	158,656	1,453.8	68,842	68,715	685.9	285	286	235.2	151,928	151,037	190.1	12,560	12,462	1,379.6	397,793	395,668	371.0			
U.S. dependent areas																											
American Samoa	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—
Guam	0	0	—	17	17	—	1	1	—	3	3	—	2	2	—	4	4	—	3	3	—	30	30	—	—	—	—
Northern Mariana Islands	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—
Puerto Rico	1	1	—	1	1	—	2	2	—	8,598	8,556	—	1	1	—	6	6	—	0	0	—	8,609	8,567	—	—	—	—
Republic of Palau	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—	0	0	—	0	0	—	1	1	—	—	—	—
U.S. Virgin Islands	0	0	—	0	0	—	156	156	—	100	101	—	0	0	—	29	29	—	1	1	—	286	286	—	—	—	—
Subtotal	1	1	—	18	18	—	159	159	—	8,701	8,659	—	4	4	—	39	39	—	4	4	—	8,926	8,883	—	—	—	—
Total^e	1,025	1,016	—	3,498	3,514	—	159,833	158,815	—	77,543	77,374	—	289	290	—	151,967	151,076	—	12,564	12,465	—	406,719	404,551	—	—	—	—

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

^d Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided for U.S. dependent areas because U.S. census information is limited for U.S. dependent areas.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 11. Adults aged 50 years and older living with diagnosed HIV infection, by population of area of residence, age group, and selected characteristics, year-end 2013—United States

	Metropolitan statistical areas (population ≥500,000)			Metropolitan areas (population of 50,000–499,999)			Nonmetropolitan areas		
	Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
50–64 years of age at end of year									
Race/ethnicity									
American Indian/Alaska Native	554	550	325.4	134	132	151.6	215	213	126.6
Asian ^b	2,680	2,696	104.8	177	178	81.6	56	56	74.8
Black/African American	116,731	116,301	2,177.0	13,085	12,988	1,276.4	7,912	7,827	1,098.2
Hispanic/Latino ^c	53,350	53,316	984.9	3,837	3,820	464.4	1,850	1,842	430.3
Native Hawaiian/Other Pacific Islander	195	196	317.4	31	31	261.0	25	25	284.7
White	106,574	106,213	392.2	14,580	14,458	163.5	7,620	7,549	90.8
Multiple races	9,480	9,429	2,319.1	1,016	1,006	1,011.0	564	559	616.5
Males									
Transmission category									
Male-to-male sexual contact	125,205	142,643	—	12,521	14,571	—	5,929	7,239	—
Injection drug use	31,635	37,598	—	3,976	4,658	—	2,782	3,276	—
Male-to-male sexual contact and injection drug use	15,924	17,294	—	2,215	2,429	—	1,371	1,523	—
Heterosexual contact ^d	19,523	25,008	—	2,660	3,449	—	1,465	1,991	—
Other ^e	31,717	716	—	4,087	145	—	2,724	108	—
Subtotal, males	224,004	223,259	1,127.2	25,459	25,253	469.0	14,271	14,137	291.0
Females									
Transmission category									
Injection drug use	16,631	21,149	—	1,712	2,090	—	852	1,081	—
Heterosexual contact ^d	31,889	43,798	—	3,758	5,201	—	1,955	2,807	—
Other ^e	17,040	495	—	1,931	70	—	1,164	44	—
Subtotal, females	65,560	65,442	308.1	7,401	7,361	128.9	3,971	3,933	79.7
Subtotal, 50–64 years of age at end of year	289,564	288,701	703.4	32,860	32,614	293.9	18,242	18,070	184.5
≥65 years of age at end of year									
Race/ethnicity									
American Indian/Alaska Native	60	59	71.7	16	16	35.4	34	34	36.7
Asian ^b	495	495	31.6	38	38	30.0	7	7	14.6
Black/African American	17,838	17,503	609.3	1,749	1,701	311.0	1,001	968	229.8
Hispanic/Latino ^c	8,456	8,395	310.0	516	511	123.0	226	225	97.0
Native Hawaiian/Other Pacific Islander	26	26	89.7	2	2	35.8	3	3	67.4
White	17,928	17,668	85.2	2,326	2,284	31.1	1,324	1,296	18.2
Multiple races	1,188	1,161	594.9	109	106	199.9	71	69	118.9
Males									
Transmission category									
Male-to-male sexual contact	18,873	21,521	—	1,801	2,125	—	956	1,146	—
Injection drug use	4,792	6,453	—	464	633	—	293	384	—
Male-to-male sexual contact and injection drug use	1,407	1,603	—	173	204	—	92	108	—
Heterosexual contact ^d	3,935	5,534	—	448	660	—	228	347	—
Other ^e	6,907	225	—	868	53	—	492	23	—
Subtotal, males	35,914	35,337	289.6	3,754	3,675	97.0	2,061	2,008	55.6
Females									
Transmission category									
Injection drug use	1,338	2,255	—	95	159	—	48	97	—
Heterosexual contact ^d	5,457	7,472	—	575	789	—	329	466	—
Other ^e	3,282	243	—	332	34	—	228	30	—
Subtotal, females	10,077	9,970	62.3	1,002	981	20.7	605	593	13.5
Subtotal, ≥65 years of age at end of year	45,991	45,307	160.7	4,756	4,657	54.6	2,666	2,601	32.5

Table 11. Adults aged 50 years and older living with diagnosed HIV infection, by population of area of residence, age group, and selected characteristics, year-end 2013—United States (cont)

	Metropolitan statistical areas (population ≥500,000)			Metropolitan areas (population of 50,000–499,999)			Nonmetropolitan areas		
	Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Total, ≥50 years of age at end of year									
Age (yr)									
50-54	614	609	242.7	150	148	112.5	249	247	94.8
55-59	3,175	3,190	77.1	215	215	62.7	63	63	51.2
60-64	134,569	133,804	1,628.8	14,834	14,689	938.9	8,913	8,795	775.7
65-69	61,806	61,711	759.9	4,353	4,331	349.9	2,076	2,066	313.2
70-74	221	222	244.0	33	33	189.4	28	28	212.6
75-79	124,502	123,881	259.0	16,906	16,742	103.5	8,944	8,844	57.3
80-84	10,668	10,590	1,759.9	1,125	1,112	729.1	635	628	421.9
≥85									
Race/ethnicity									
American Indian/Alaska Native	96,629	96,249	686.0	11,017	10,924	286.9	6,131	6,070	180.1
Asian ^b	56,885	56,459	481.4	6,044	5,963	177.3	3,390	3,334	109.6
Black/African American	26,970	26,658	287.2	2,853	2,806	101.2	1,583	1,553	60.7
Hispanic/Latino ^c	11,722	11,542	174.6	1,220	1,194	58.3	686	670	34.3
Native Hawaiian/Other Pacific Islander	4,730	4,629	97.0	438	424	28.4	275	266	18.7
White	1,827	1,773	48.7	162	154	13.9	81	75	7.3
Multiple races	742	704	18.1	83	78	7.0	41	37	3.6
Males									
Transmission category									
Male-to-male sexual contact	144,078	164,164	—	14,322	16,696	—	6,885	8,385	—
Injection drug use	36,427	44,051	—	4,440	5,290	—	3,075	3,660	—
Male-to-male sexual contact and injection drug use	17,331	18,896	—	2,388	2,633	—	1,463	1,631	—
Heterosexual contact ^d	23,458	30,542	—	3,108	4,109	—	1,693	2,338	—
Other ^e	38,624	941	—	4,955	199	—	3,216	131	—
Subtotal, males	259,918	258,595	807.9	29,213	28,928	315.3	16,332	16,146	190.6
Females									
Transmission category									
Injection drug use	17,969	23,404	—	1,807	2,249	—	900	1,179	—
Heterosexual contact ^d	37,346	51,270	—	4,333	5,990	—	2,284	3,273	—
Other ^e	20,322	738	—	2,263	104	—	1,392	74	—
Subtotal, females	75,637	75,412	202.5	8,403	8,343	79.8	4,576	4,526	48.6
Total^f	335,555	334,008	482.4	37,616	37,270	189.9	20,908	20,672	116.2

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 12a. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
Males												
Age at end of year (yr)												
50–54	2,397	2,508	22.9	2,372	2,525	22.8	2,301	2,499	22.5	2,054	2,371	21.4
55–59	1,961	2,051	21.4	2,032	2,163	22.0	1,999	2,168	21.5	1,781	2,058	20.0
60–64	1,341	1,403	17.2	1,382	1,471	17.2	1,442	1,569	18.4	1,319	1,522	17.5
65–69	686	719	12.2	739	790	13.0	798	865	13.1	797	927	13.4
70–74	365	382	8.9	379	404	9.2	428	465	10.1	414	481	9.9
75–79	192	202	6.3	199	212	6.6	214	231	7.0	205	234	6.9
80–84	77	81	3.5	75	80	3.4	95	104	4.4	105	122	5.1
≥85	39	41	2.3	31	33	1.7	44	48	2.5	43	51	2.5
Race/ethnicity												
American Indian/Alaska Native	18	19	7.0	19	20	7.1	16	18	6.1	21	24	8.0
Asian ^b	29	30	1.7	38	41	2.2	29	31	1.6	26	29	1.4
Black/African American	3,097	3,240	74.5	3,198	3,408	75.7	3,140	3,418	73.6	2,754	3,221	67.3
Hispanic/Latino ^c	1,046	1,087	27.3	1,092	1,151	27.3	1,065	1,142	25.7	1,085	1,225	26.1
Native Hawaiian/Other Pacific Islander	2	2	4.1	5	5	10.0	4	4	7.3	2	3	4.3
White	2,508	2,636	7.4	2,534	2,710	7.5	2,685	2,924	8.0	2,500	2,888	7.7
Multiple races	358	374	101.8	323	342	88.9	382	412	102.5	330	376	89.7
Transmission category												
Male-to-male sexual contact	2,671	3,401	—	2,694	3,522	—	2,861	3,792	—	2,798	3,845	—
Injection drug use	1,790	2,152	—	1,760	2,165	—	1,725	2,156	—	1,486	1,985	—
Male-to-male sexual contact and injection drug use	599	681	—	674	779	—	632	748	—	633	784	—
Heterosexual contact ^d	753	1,102	—	782	1,163	—	804	1,199	—	702	1,107	—
Other ^e	1,245	51	—	1,299	48	—	1,299	54	—	1,099	44	—
Subtotal, males	7,058	7,387	16.0	7,209	7,677	16.2	7,321	7,949	16.4	6,718	7,765	15.6
Females												
Age at end of year (yr)												
50–54	784	819	7.2	772	819	7.1	804	870	7.6	654	756	6.6
55–59	602	632	6.2	619	656	6.3	641	694	6.5	605	700	6.4
60–64	348	364	4.1	345	366	3.9	426	463	5.0	396	464	4.9
65–69	167	175	2.6	197	209	3.1	187	204	2.8	203	231	3.0
70–74	96	101	2.0	101	108	2.1	96	105	1.9	99	120	2.1
75–79	30	32	0.8	59	63	1.5	55	60	1.4	51	59	1.4
80–84	27	28	0.8	28	30	0.9	28	31	0.9	25	30	0.9
≥85	11	12	0.3	17	18	0.5	28	31	0.8	8	10	0.2
Race/ethnicity												
American Indian/Alaska Native	7	7	2.4	7	7	2.3	7	8	2.3	7	8	2.2
Asian ^b	10	10	0.5	7	7	0.3	6	6	0.3	5	7	0.3
Black/African American	1,312	1,374	24.5	1,311	1,392	24.1	1,352	1,469	24.7	1,222	1,427	23.3
Hispanic/Latino ^c	276	286	6.3	325	342	7.1	328	352	7.0	301	342	6.4
Native Hawaiian/Other Pacific Islander	0	0	0.0	1	1	1.8	0	0	0.0	0	0	0.0
White	361	381	0.9	396	423	1.0	446	486	1.2	384	447	1.1
Multiple races	99	104	24.4	91	97	21.8	126	137	29.5	122	139	28.8
Transmission category												
Injection drug use	696	883	—	752	935	—	777	1,012	—	725	992	—
Heterosexual contact ^d	845	1,259	—	864	1,303	—	899	1,422	—	838	1,350	—
Other ^e	524	20	—	522	31	—	589	26	—	478	27	—
Subtotal, females	2,065	2,162	4.0	2,138	2,269	4.2	2,265	2,459	4.4	2,041	2,369	4.2

Table 12a. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Both sexes												
Age at end of year (yr)												
50–54	3,181	3,327	14.9	3,144	3,345	14.8	3,105	3,370	14.9	2,708	3,127	13.8
55–59	2,563	2,682	13.6	2,651	2,818	13.9	2,640	2,862	13.8	2,386	2,757	13.0
60–64	1,689	1,767	10.4	1,727	1,837	10.3	1,868	2,032	11.4	1,715	1,985	10.9
65–69	853	894	7.1	936	999	7.8	985	1,069	7.6	1,000	1,159	7.9
70–74	461	483	5.2	480	511	5.3	524	570	5.7	513	601	5.7
75–79	222	234	3.2	258	275	3.7	269	292	3.9	256	293	3.8
80–84	104	109	1.9	103	110	1.9	123	135	2.3	130	151	2.6
≥85	50	53	1.0	48	51	0.9	72	79	1.3	51	61	1.0
Race/ethnicity												
American Indian/Alaska Native	25	26	4.5	26	27	4.6	23	25	4.1	28	32	4.9
Asian ^b	39	41	1.0	45	48	1.1	35	38	0.9	31	36	0.8
Black/African American	4,409	4,614	46.4	4,509	4,800	46.7	4,492	4,888	46.1	3,976	4,648	42.6
Hispanic/Latino ^c	1,322	1,373	16.1	1,417	1,493	16.5	1,393	1,494	15.7	1,386	1,567	15.6
Native Hawaiian/Other Pacific Islander	2	2	2.0	6	6	5.8	4	4	3.5	2	3	2.1
White	2,869	3,017	4.0	2,930	3,133	4.1	3,131	3,410	4.4	2,884	3,335	4.2
Multiple races	457	478	60.2	414	439	52.9	508	549	63.4	452	515	57.0
Total^f	9,123	9,550	9.6	9,347	9,946	9.7	9,586	10,408	10.0	8,759	10,134	9.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 12b. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
Males												
Age at end of year (yr)												
50–54	2,458	2,572	23.2	2,452	2,612	23.3	2,354	2,558	22.8	2,113	2,440	21.8
55–59	2,007	2,099	21.6	2,087	2,222	22.4	2,049	2,224	21.8	1,824	2,108	20.3
60–64	1,378	1,442	17.4	1,420	1,512	17.5	1,473	1,603	18.5	1,355	1,563	17.8
65–69	707	742	12.4	759	811	13.2	821	891	13.3	818	952	13.6
70–74	373	390	9.0	391	417	9.3	439	477	10.2	422	491	9.9
75–79	202	213	6.6	207	221	6.7	226	245	7.3	211	241	7.0
80–84	80	84	3.6	79	84	3.6	99	108	4.5	109	126	5.3
≥85	43	45	2.5	33	35	1.8	44	48	2.4	44	52	2.5
Race/ethnicity												
American Indian/Alaska Native	18	19	—	19	20	—	16	18	—	21	24	—
Asian ^b	29	30	—	38	41	—	29	31	—	26	29	—
Black/African American	3,099	3,242	—	3,202	3,412	—	3,145	3,424	—	2,756	3,223	—
Hispanic/Latino ^c	1,232	1,283	—	1,307	1,383	—	1,243	1,340	—	1,259	1,428	—
Native Hawaiian/Other Pacific Islander	2	2	—	5	5	—	4	4	—	2	3	—
White	2,510	2,638	—	2,534	2,710	—	2,685	2,924	—	2,500	2,888	—
Multiple races	358	374	—	323	342	—	383	413	—	332	378	—
Transmission category												
Male-to-male sexual contact	2,702	3,436	—	2,740	3,572	—	2,887	3,822	—	2,838	3,893	—
Injection drug use	1,873	2,244	—	1,861	2,278	—	1,803	2,249	—	1,552	2,065	—
Male-to-male sexual contact and injection drug use	614	697	—	685	791	—	644	761	—	643	795	—
Heterosexual contact ^d	804	1,159	—	835	1,224	—	862	1,266	—	758	1,176	—
Other ^e	1,255	52	—	1,307	48	—	1,309	55	—	1,105	44	—
Subtotal, males	7,248	7,587	16.2	7,428	7,914	16.5	7,505	8,154	16.6	6,896	7,974	15.9
Females												
Age at end of year (yr)												
50–54	810	847	7.3	799	849	7.3	825	893	7.7	675	780	6.7
55–59	620	651	6.3	634	672	6.3	657	712	6.6	621	719	6.5
60–64	359	375	4.2	355	377	4.0	437	476	5.1	406	475	5.0
65–69	178	186	2.8	202	214	3.1	191	209	2.8	213	243	3.1
70–74	97	102	2.0	106	113	2.1	100	110	2.0	101	122	2.1
75–79	31	33	0.8	61	65	1.5	57	62	1.5	54	62	1.4
80–84	27	28	0.8	30	33	0.9	30	33	1.0	27	33	0.9
≥85	11	12	0.3	18	19	0.5	30	33	0.8	9	11	0.3
Race/ethnicity												
American Indian/Alaska Native	7	7	—	7	7	—	7	8	—	7	8	—
Asian ^b	10	10	—	7	7	—	6	6	—	5	7	—
Black/African American	1,312	1,374	—	1,311	1,392	—	1,353	1,470	—	1,224	1,430	—
Hispanic/Latino ^c	344	358	—	391	414	—	389	420	—	364	415	—
Native Hawaiian/Other Pacific Islander	0	0	—	1	1	—	0	0	—	0	0	—
White	361	381	—	397	424	—	446	486	—	384	447	—
Multiple races	99	104	—	91	97	—	126	137	—	122	139	—
Transmission category												
Injection drug use	718	907	—	766	950	—	791	1,028	—	742	1,013	—
Heterosexual contact ^d	891	1,307	—	916	1,361	—	942	1,473	—	882	1,405	—
Other ^e	524	20	—	523	31	—	594	27	—	482	27	—
Subtotal, females	2,133	2,234	4.1	2,205	2,342	4.2	2,327	2,528	4.5	2,106	2,445	4.2

Table 12b. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year, sex, and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Both sexes												
Age at end of year (yr)												
50–54	3,268	3,419	15.1	3,251	3,460	15.1	3,179	3,451	15.1	2,788	3,221	14.1
55–59	2,627	2,750	13.7	2,721	2,894	14.1	2,706	2,936	14.0	2,445	2,827	13.2
60–64	1,737	1,818	10.6	1,775	1,889	10.5	1,910	2,079	11.5	1,761	2,038	11.1
65–69	885	928	7.3	961	1,026	7.8	1,012	1,099	7.7	1,031	1,195	8.1
70–74	470	493	5.2	497	530	5.4	539	587	5.8	523	613	5.7
75–79	233	246	3.3	268	286	3.8	283	307	4.0	265	304	3.9
80–84	107	112	1.9	109	117	2.0	129	141	2.4	136	159	2.7
≥85	54	57	1.0	51	54	0.9	74	81	1.4	53	63	1.0
Race/ethnicity												
American Indian/Alaska Native	25	26	—	26	27	—	23	25	—	28	32	—
Asian ^b	39	41	—	45	48	—	35	38	—	31	36	—
Black/African American	4,411	4,616	—	4,513	4,804	—	4,498	4,894	—	3,980	4,653	—
Hispanic/Latino ^c	1,576	1,641	—	1,698	1,797	—	1,632	1,760	—	1,623	1,844	—
Native Hawaiian/Other Pacific Islander	2	2	—	6	6	—	4	4	—	2	3	—
White	2,871	3,019	—	2,931	3,134	—	3,131	3,410	—	2,884	3,335	—
Multiple races	457	478	—	414	439	—	509	550	—	454	518	—
Total^f	9,381	9,822	9.7	9,633	10,256	9.9	9,832	10,682	10.1	9,002	10,419	9.6

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 13a. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
50–64 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	16	17	8.9	15	16	8.1	15	16	8.3	17	20	9.8
Asian ^b	21	22	1.9	30	32	2.6	19	21	1.6	18	20	1.6
Black/African American	2,520	2,635	87.6	2,585	2,753	88.4	2,471	2,690	84.4	2,104	2,462	75.6
Hispanic/Latino ^c	842	874	31.4	864	911	31.0	851	913	29.5	863	973	29.9
Native Hawaiian/Other Pacific Islander	2	2	5.7	4	4	11.5	3	3	7.8	2	3	6.2
White	2,001	2,102	9.9	2,021	2,160	10.0	2,063	2,246	10.4	1,881	2,167	10.0
Multiple races	297	310	121.2	267	282	105.5	320	346	125.6	269	306	107.5
Transmission category												
Male-to-male sexual contact	2,110	2,681	—	2,128	2,772	—	2,218	2,930	—	2,110	2,902	—
Injection drug use	1,589	1,856	—	1,542	1,835	—	1,489	1,795	—	1,238	1,601	—
Male-to-male sexual contact and injection drug use	546	616	—	623	711	—	560	657	—	574	700	—
Heterosexual contact ^d	554	787	—	565	813	—	581	827	—	468	725	—
Other ^e	900	23	—	928	28	—	894	28	—	764	22	—
Subtotal, males	5,699	5,962	20.8	5,786	6,159	20.9	5,742	6,236	21.0	5,154	5,951	19.8
Females												
Race/ethnicity												
American Indian/Alaska Native	7	7	3.5	4	4	2.0	6	6	3.0	6	7	3.0
Asian ^b	6	6	0.4	3	3	0.2	3	3	0.2	4	6	0.4
Black/African American	1,103	1,154	32.7	1,072	1,138	31.2	1,124	1,220	32.6	984	1,147	30.1
Hispanic/Latino ^c	226	234	7.9	251	263	8.5	265	284	8.7	231	263	7.7
Native Hawaiian/Other Pacific Islander	0	0	0	1	1	2.6	0	0	0.0	0	0	0.0
White	311	328	1.5	329	350	1.6	368	401	1.8	331	385	1.7
Multiple races	81	85	30.4	76	81	27.6	105	114	37.7	99	112	35.9
Transmission category												
Injection drug use	656	802	—	691	826	—	725	905	—	655	865	—
Heterosexual contact ^d	672	1003	—	673	999	—	711	1,107	—	651	1,040	—
Other ^e	406	10	—	372	16	—	435	16	—	349	14	—
Subtotal, females	1,734	1,815	6	1,736	1,841	5.9	1,871	2,028	6.4	1,655	1,919	6.0
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	23	24	6.1	19	20	4.9	21	23	5.5	23	26	6.2
Asian ^b	27	28	1.1	33	35	1.3	22	24	0.9	22	26	0.9
Black/African American	3,623	3,789	58.0	3,657	3,891	57.5	3,595	3,910	56.5	3,088	3,609	51.0
Hispanic/Latino ^c	1,068	1,109	19.3	1,115	1,175	19.4	1,116	1,197	18.8	1,094	1,237	18.6
Native Hawaiian/Other Pacific Islander	2	2	2.8	5	5	7.0	3	3	3.8	2	3	3.1
White	2,312	2,430	5.6	2,350	2,510	5.7	2,431	2,647	6.0	2,212	2,552	5.8
Multiple races	378	395	73.8	343	363	64.8	425	460	79.6	368	418	70.0
Subtotal, 50–64 years of age at end of year	7,433	7,776	13.1	7,522	7,999	13.2	7,613	8,264	13.5	6,809	7,870	12.7

Table 13a. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
≥65 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	2	2	2.6	4	4	4.8	1	1	1.3	4	4	4.3
Asian ^b	8	8	1.4	8	9	1.4	10	11	1.6	8	9	1.2
Black/African American	577	605	45.1	613	655	47.3	669	728	49.9	650	759	49.6
Hispanic/Latino ^c	204	212	17.7	228	240	18.9	214	228	16.9	222	251	17.4
Native Hawaiian/Other Pacific Islander	0	0	0.0	1	1	6.5	1	1	6.1	0	0	0.0
White	507	534	3.8	513	550	3.8	622	678	4.5	619	721	4.6
Multiple races	61	64	57.4	56	60	51.1	62	66	52.1	61	70	52.1
Transmission category												
Male-to-male sexual contact	561	721	—	566	750	—	643	862	—	688	944	—
Injection drug use	201	296	—	218	329	—	236	361	—	248	384	—
Male-to-male sexual contact and injection drug use	53	65	—	51	68	—	72	92	—	59	83	—
Heterosexual contact ^d	199	315	—	217	350	—	223	373	—	234	382	—
Other ^e	345	28	—	371	21	—	405	26	—	335	22	—
Subtotal, males	1,359	1,425	8.2	1,423	1,519	8.5	1,579	1,713	9.1	1,564	1,815	9.3
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	0.0	3	3	3.1	1	1	1.1	1	1	0.9
Asian ^b	4	4	0.5	4	4	0.5	3	3	0.4	1	1	0.1
Black/African American	209	220	10.6	239	253	11.9	228	249	11.2	238	280	12.1
Hispanic/Latino ^c	50	52	3.2	74	78	4.6	63	68	3.8	70	79	4.1
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	50	53	0.3	67	73	0.4	78	86	0.4	53	62	0.3
Multiple races	18	19	12.8	15	16	10.7	21	23	14.2	23	27	15.8
Transmission category												
Injection drug use	40	82	—	61	109	—	52	106	—	70	128	—
Heterosexual contact ^d	173	256	—	191	304	—	188	315	—	187	310	—
Other ^e	118	10	—	150	15	—	154	10	—	129	12	—
Subtotal, females	331	348	1.5	402	428	1.8	394	431	1.8	386	450	1.8
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	2	2	1.2	7	7	3.8	2	2	1.2	5	5	2.4
Asian ^b	12	13	0.9	12	13	0.9	13	14	0.9	9	10	0.6
Black/African American	786	825	24.2	852	909	25.9	897	977	26.6	888	1,039	27.1
Hispanic/Latino ^c	254	264	9.4	302	318	10.7	277	297	9.4	292	330	9.8
Native Hawaiian/Other Pacific Islander	0	0	0.0	1	1	3.0	1	1	2.8	0	0	0.0
White	557	587	1.8	580	623	1.9	700	764	2.2	672	783	2.2
Multiple races	79	83	32.1	71	76	28.3	83	89	30.8	84	97	31.8
Subtotal, ≥65 years of age at end of year	1,690	1,773	4.4	1,825	1,947	4.7	1,973	2,144	5.0	1,950	2,265	5.1
Total^f	9,123	9,550	9.6	9,347	9,946	9.7	9,586	10,408	10.0	8,759	10,134	9.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 13b. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States and 6 dependent areas

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate									
50–64 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	16	17	—	15	16	—	15	16	—	17	20	—
Asian ^b	21	22	—	30	32	—	19	21	—	18	20	—
Black/African American	2,521	2,636	—	2,588	2,756	—	2,473	2,693	—	2,105	2,463	—
Hispanic/Latino ^c	985	1,025	—	1,034	1,096	—	982	1,059	—	998	1,131	—
Native Hawaiian/Other Pacific Islander	2	2	—	4	4	—	3	3	—	2	3	—
White	2,001	2,102	—	2,021	2,160	—	2,063	2,246	—	1,881	2,167	—
Multiple races	297	310	—	267	282	—	321	347	—	271	308	—
Transmission category												
Male-to-male sexual contact	2,131	2,704	—	2,165	2,812	—	2,235	2,950	—	2,139	2,936	—
Injection drug use	1,664	1,938	—	1,627	1,931	—	1,557	1,875	—	1,295	1,669	—
Male-to-male sexual contact and injection drug use	560	630	—	633	722	—	570	667	—	584	711	—
Heterosexual contact ^d	581	817	—	599	853	—	614	864	—	506	773	—
Other ^e	907	24	—	935	28	—	900	28	—	768	22	—
Subtotal, males	5,843	6,113	21.0	5,959	6,346	21.3	5,876	6,385	21.3	5,292	6,112	20.1
Females												
Race/ethnicity												
American Indian/Alaska Native	7	7	—	4	4	—	6	6	—	6	7	—
Asian ^b	6	6	—	3	3	—	3	3	—	4	6	—
Black/African American	1,103	1,154	—	1,072	1,138	—	1,125	1,221	—	985	1,148	—
Hispanic/Latino ^c	281	293	—	302	319	—	312	336	—	277	316	—
Native Hawaiian/Other Pacific Islander	0	0	—	1	1	—	0	0	—	0	0	—
White	311	328	—	330	351	—	368	401	—	331	385	—
Multiple races	81	85	—	76	81	—	105	114	—	99	112	—
Transmission category												
Injection drug use	676	823	—	704	840	—	738	921	—	670	883	—
Heterosexual contact ^d	707	1,040	—	712	1,042	—	742	1,145	—	681	1,077	—
Other ^e	406	10	—	372	16	—	439	16	—	351	14	—
Subtotal, females	1,789	1,873	6.1	1,788	1,898	6.0	1,919	2,081	6.5	1,702	1,974	6.1
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	23	24	—	19	20	—	21	23	—	23	26	—
Asian ^b	27	28	—	33	35	—	22	24	—	22	26	—
Black/African American	3,624	3,790	—	3,660	3,895	—	3,598	3,914	—	3,090	3,611	—
Hispanic/Latino ^c	1,266	1,318	—	1,336	1,415	—	1,294	1,395	—	1,275	1,447	—
Native Hawaiian/Other Pacific Islander	2	2	—	5	5	—	3	3	—	2	3	—
White	2,312	2,430	—	2,351	2,512	—	2,431	2,647	—	2,212	2,552	—
Multiple races	378	395	—	343	363	—	426	461	—	370	420	—
Subtotal, 50–64 years of age at end of year	7,632	7,986	13.3	7,747	8,244	13.4	7,795	8,466	13.7	6,994	8,085	12.9

Table 13b. Deaths of adults aged 50 years and older living with diagnosed HIV infection, by year of death, age at death, and selected characteristics, 2010–2013—United States and 6 dependent areas (cont)

	2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
≥65 years of age at end of year												
Males												
Race/ethnicity												
American Indian/Alaska Native	2	2	—	4	4	—	1	1	—	4	4	—
Asian ^b	8	8	—	8	9	—	10	11	—	8	9	—
Black/African American	578	606	—	614	656	—	672	731	—	651	760	—
Hispanic/Latino ^c	247	257	—	273	288	—	261	281	—	261	297	—
Native Hawaiian/Other Pacific Islander	0	0	—	1	1	—	1	1	—	0	0	—
White	509	536	—	513	550	—	622	678	—	619	721	—
Multiple races	61	64	—	56	60	—	62	66	—	61	70	—
Transmission category												
Male-to-male sexual contact	571	732	—	575	760	—	652	872	—	699	958	—
Injection drug use	209	306	—	234	347	—	246	374	—	257	396	—
Male-to-male sexual contact and injection drug use	54	66	—	52	69	—	74	94	—	59	83	—
Heterosexual contact ^d	223	342	—	236	371	—	248	402	—	252	403	—
Other ^e	348	28	—	372	21	—	409	27	—	337	22	—
Subtotal, males	1,405	1,474	8.3	1,469	1,568	8.6	1,629	1,769	9.3	1,604	1,862	9.4
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	—	3	3	—	1	1	—	1	1	—
Asian ^b	4	4	—	4	4	—	3	3	—	1	1	—
Black/African American	209	220	—	239	253	—	228	249	—	239	282	—
Hispanic/Latino ^c	63	65	—	89	95	—	77	84	—	87	99	—
Native Hawaiian/Other Pacific Islander	50	53	—	67	73	—	78	86	—	53	62	—
White	18	19	—	15	16	—	21	23	—	23	27	—
Multiple races			—			—			—			—
Transmission category												
Injection drug use	42	84	—	62	111	—	53	107	—	72	130	—
Heterosexual contact ^d	184	267	—	204	319	—	200	328	—	201	329	—
Other ^e	118	10	—	151	15	—	155	11	—	131	12	—
Subtotal, females	344	361	1.5	417	445	1.9	408	447	1.8	404	471	1.9
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	2	2	—	7	7	—	2	2	—	5	5	—
Asian ^b	12	13	—	12	13	—	13	14	—	9	10	—
Black/African American	787	826	—	853	910	—	900	981	—	890	1,042	—
Hispanic/Latino ^c	310	323	—	362	382	—	338	365	—	348	396	—
Native Hawaiian/Other Pacific Islander	0	0	—	1	1	—	1	1	—	0	0	—
White	559	589	—	580	623	—	700	764	—	672	783	—
Multiple races	79	83	—	71	76	—	83	89	—	84	97	—
Subtotal, ≥65 years of age at end of year	1,749	1,835	4.5	1,886	2,012	4.8	2,037	2,215	5.1	2,008	2,333	5.1
Total^f	9,381	9,822	9.7	9,633	10,256	9.9	9,832	10,682	10.1	9,002	10,419	9.6

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 14a. Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States

	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate									
Males												
Race/ethnicity												
American Indian/Alaska Native	9	11	14.4	6	6	9.3	2	2	4.0	3	3	7.8
Asian ^b	8	9	1.8	5	5	1.2	5	6	1.6	4	5	1.7
Black/African American	768	897	70.4	767	895	79.7	569	670	77.9	339	399	67.4
Hispanic/Latino ^c	380	432	30.7	269	302	28.1	214	240	31.0	114	129	23.7
Native Hawaiian/Other Pacific Islander	2	3	15.2	0	0	0.0	0	0	0.0	0	0	0.0
White	767	883	11.5	651	754	10.1	463	529	8.1	312	362	6.7
Multiple races	120	136	120.3	83	95	97.0	66	75	101.6	25	29	54.3
Transmission category												
Male-to-male sexual contact	922	1,256	—	692	955	—	496	691	—	345	470	—
Injection drug use	388	518	—	470	589	—	380	493	—	165	231	—
Male-to-male sexual contact and injection drug use	248	306	—	208	251	—	118	144	—	40	55	—
Heterosexual contact ^d	199	285	—	166	256	—	103	184	—	102	164	—
Other ^e	297	7	—	245	6	—	222	10	—	145	6	—
Subtotal, males	2,054	2,371	21.4	1,781	2,058	20.0	1,319	1,522	17.5	797	927	13.4
Females												
Race/ethnicity												
American Indian/Alaska Native	4	4	4.9	2	2	3.1	0	0	0.0	1	1	2.3
Asian ^b	0	0	0.0	3	4	0.8	1	1	0.3	0	0	0.0
Black/African American	381	442	30.6	367	425	32.4	236	279	26.3	124	142	18.6
Hispanic/Latino ^c	89	101	7.2	73	83	7.3	69	79	9.2	36	40	6.2
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	133	155	2.0	129	150	1.9	69	81	1.2	31	36	0.6
Multiple races	47	54	43.1	31	35	32.9	21	24	28.9	11	13	21.5
Transmission category												
Injection drug use	263	346	—	246	322	—	146	196	—	53	80	—
Heterosexual contact ^d	254	402	—	230	373	—	167	265	—	94	151	—
Other ^e	137	7	—	129	5	—	83	3	—	56	1	—
Subtotal, females	654	756	6.6	605	700	6.4	396	464	4.9	203	231	3.0
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	13	16	9.5	8	9	6.0	2	2	1.9	4	4	4.9
Asian ^b	8	9	0.9	8	10	1.0	6	7	0.9	4	5	0.8
Black/African American	1,149	1,339	49.2	1,134	1,321	54.3	805	949	49.4	463	541	39.9
Hispanic/Latino ^c	469	533	18.9	342	385	17.4	283	319	19.5	150	169	14.2
Native Hawaiian/Other Pacific Islander	2	3	7.6	0	0	0.0	0	0	0.0	0	0	0.0
White	900	1,038	6.7	780	904	5.9	532	610	4.5	343	398	3.5
Multiple races	167	190	79.9	114	130	63.6	87	98	63.4	36	42	36.8
Total^f	2,708	3,127	13.8	2,386	2,757	13.0	1,715	1,985	10.9	1,000	1,159	7.9

Table 14a. Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States (cont)

	70–74 years			75–79 years			80–84 years			≥85 years		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	1	1	4.2	0	0	0.0	0	0	0.0	0	0	0.0
Asian ^b	3	3	1.6	0	0	0.0	1	1	1.3	0	0	0.0
Black/African American	169	197	49.9	81	93	35.3	40	45	28.8	21	25	19.9
Hispanic/Latino ^c	57	64	17.8	29	32	12.9	14	16	10.2	8	10	7.6
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	166	195	5.0	84	97	3.6	45	53	2.7	12	14	0.8
Multiple races	18	21	60.3	11	12	55.3	5	6	43.2	2	2	19.6
Transmission category												
Male-to-male sexual contact	188	262	—	90	121	—	48	66	—	17	25	—
Injection drug use	54	91	—	19	40	—	7	14	—	3	7	—
Male-to-male sexual contact and injection drug use	13	19	—	3	5	—	3	4	—	0	1	—
Heterosexual contact ^d	60	104	—	45	64	—	21	34	—	6	15	—
Other ^e	99	6	—	48	4	—	26	4	—	17	2	—
Subtotal, males	414	481	9.9	205	234	6.9	105	122	5.1	43	51	2.5
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
Asian ^b	0	0	0.0	0	0	0.0	1	1	0.8	0	0	0.0
Black/African American	66	81	14.7	30	36	8.8	12	14	5.0	6	8	2.5
Hispanic/Latino ^c	17	19	4.2	10	11	3.2	7	9	3.6	0	0	0.0
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	10	12	0.3	9	10	0.3	2	2	0.1	1	1	0.0
Multiple races	6	7	17.5	2	2	7.6	3	4	17.7	1	1	5.3
Transmission category												
Injection drug use	13	30	—	2	10	—	2	7	—	0	1	—
Heterosexual contact ^d	50	87	—	26	46	—	12	19	—	5	7	—
Other ^e	36	3	—	23	3	—	11	4	—	3	1	—
Subtotal, females	99	120	2.1	51	59	1.4	25	30	0.9	8	10	0.2
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	1	1	2.0	0	0	0.0	0	0	0.0	0	0	0.0
Asian ^b	3	3	0.7	0	0	0.0	2	2	1.0	0	0	0.0
Black/African American	235	277	29.4	111	128	19.2	52	60	13.4	27	33	7.6
Hispanic/Latino ^c	74	83	10.2	39	43	7.3	21	25	6.2	8	10	2.7
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0
White	176	208	2.5	93	107	1.8	47	55	1.2	13	15	0.3
Multiple races	24	28	37.0	13	15	28.6	8	10	28.0	3	3	10.2
Total^f	513	601	5.7	256	293	3.8	130	151	2.6	51	61	1.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 14b. Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States and 6 dependent areas

	50–54 years			55–59 years			60–64 years			65–69 years		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate									
Males												
Race/ethnicity												
American Indian/Alaska Native	9	11	—	6	6	—	2	2	—	3	3	—
Asian ^b	8	9	—	5	5	—	5	6	—	4	5	—
Black/African American	769	898	—	767	895	—	569	670	—	339	399	—
Hispanic/Latino ^c	437	498	—	311	352	—	250	281	—	135	153	—
Native Hawaiian/Other Pacific Islander	2	3	—	0	0	—	0	0	—	0	0	—
White	767	883	—	651	754	—	463	529	—	312	362	—
Multiple races	121	138	—	84	96	—	66	75	—	25	29	—
Transmission category												
Male-to-male sexual contact	930	1,265	—	701	966	—	508	705	—	350	477	—
Injection drug use	420	557	—	482	604	—	393	508	—	171	239	—
Male-to-male sexual contact and injection drug use	252	310	—	211	254	—	121	147	—	40	55	—
Heterosexual contact ^d	213	302	—	183	278	—	110	193	—	111	175	—
Other ^e	298	7	—	247	6	—	223	10	—	146	6	—
Subtotal, males	2,113	2,440	21.8	1,824	2,108	20.3	1,355	1,563	17.8	818	952	13.6
Females												
Race/ethnicity												
American Indian/Alaska Native	4	4	—	2	2	—	0	0	—	1	1	—
Asian ^b	0	0	—	3	4	—	1	1	—	0	0	—
Black/African American	382	444	—	367	425	—	236	279	—	124	142	—
Hispanic/Latino ^c	109	124	—	89	102	—	79	90	—	46	51	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	0	0	—
White	133	155	—	129	150	—	69	81	—	31	36	—
Multiple races	47	54	—	31	35	—	21	24	—	11	13	—
Transmission category												
Injection drug use	270	355	—	251	328	—	149	200	—	53	80	—
Heterosexual contact ^d	266	418	—	241	386	—	174	273	—	103	163	—
Other ^e	139	7	—	129	5	—	83	3	—	57	1	—
Subtotal, females	675	780	6.7	621	719	6.5	406	475	5.0	213	243	3.1
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	13	16	—	8	9	—	2	2	—	4	4	—
Asian ^b	8	9	—	8	10	—	6	7	—	4	5	—
Black/African American	1,151	1,342	—	1,134	1,321	—	805	949	—	463	541	—
Hispanic/Latino ^c	546	622	—	400	453	—	329	372	—	181	205	—
Native Hawaiian/Other Pacific Islander	2	3	—	0	0	—	0	0	—	0	0	—
White	900	1,038	—	780	904	—	532	610	—	343	398	—
Multiple races	168	191	—	115	131	—	87	98	—	36	42	—
Total^f	2,788	3,221	14.1	2,445	2,827	13.2	1,761	2,038	11.1	1,031	1,195	8.1

Table 14b. Deaths of adults aged 50 years and older with diagnosed HIV infection, by 5-year age group (age at death) and selected characteristics, 2013—United States and 6 dependent areas (cont)

	70–74 years			75–79 years			80–84 years			≥85 years		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Males												
Race/ethnicity												
American Indian/Alaska Native	1	1	—	0	0	—	0	0	—	0	0	—
Asian ^b	3	3	—	0	0	—	1	1	—	0	0	—
Black/African American	170	198	—	81	93	—	40	45	—	21	25	—
Hispanic/Latino ^c	64	72	—	35	39	—	18	21	—	9	11	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	0	0	—
White	166	195	—	84	97	—	45	53	—	12	14	—
Multiple races	18	21	—	11	12	—	5	6	—	2	2	—
Transmission category												
Male-to-male sexual contact	190	264	—	93	125	—	48	66	—	18	27	—
Injection drug use	55	93	—	20	41	—	8	16	—	3	7	—
Male-to-male sexual contact and injection drug use	13	19	—	3	5	—	3	4	—	0	1	—
Heterosexual contact ^d	64	109	—	47	67	—	24	38	—	6	15	—
Other ^e	100	6	—	48	4	—	26	4	—	17	2	—
Subtotal, males	422	491	9.9	211	241	7.0	109	126	5.3	44	52	2.5
Females												
Race/ethnicity												
American Indian/Alaska Native	0	0	—	0	0	—	0	0	—	0	0	—
Asian ^b	0	0	—	0	0	—	1	1	—	0	0	—
Black/African American	67	82	—	30	36	—	12	14	—	6	8	—
Hispanic/Latino ^c	18	21	—	13	14	—	9	11	—	1	1	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	0	0	—
White	10	12	—	9	10	—	2	2	—	1	1	—
Multiple races	6	7	—	2	2	—	3	4	—	1	1	—
Transmission category												
Injection drug use	13	30	—	3	11	—	3	8	—	0	1	—
Heterosexual contact ^d	51	89	—	28	48	—	13	21	—	6	8	—
Other ^e	37	3	—	23	3	—	11	4	—	3	1	—
Subtotal, females	101	122	2.1	54	62	1.4	27	33	0.9	9	11	0.3
Both sexes												
Race/ethnicity												
American Indian/Alaska Native	1	1	—	0	0	—	0	0	—	0	0	—
Asian ^b	3	3	—	0	0	—	2	2	—	0	0	—
Black/African American	237	280	—	111	128	—	52	60	—	27	33	—
Hispanic/Latino ^c	82	93	—	48	54	—	27	32	—	10	12	—
Native Hawaiian/Other Pacific Islander	0	0	—	0	0	—	0	0	—	0	0	—
White	176	208	—	93	107	—	47	55	—	13	15	—
Multiple races	24	28	—	13	15	—	8	10	—	3	3	—
Total^f	523	613	5.7	265	304	3.9	136	159	2.7	53	63	1.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosis of HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Includes persons of unknown race/ethnicity. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15a. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
	Est. No. ^c	Est. No. ^c	%	Est. No. ^c	%
Males					
Age at diagnosis (yr)					
50–54	2,359	909	38.5	1,450	61.5
55–59	1,553	591	38.0	962	62.0
60–64	847	314	37.1	532	62.9
65–69	419	183	43.8	236	56.2
70–74	167	67	40.2	100	59.8
75–79	64	29	45.9	35	54.1
80–84	28	10	35.3	18	64.7
≥85	12	1	8.6	11	91.4
Race/ethnicity					
American Indian/Alaska Native	16	6	40.5	9	59.5
Asian	60	35	58.5	25	41.5
Black/African American	1,951	700	35.9	1,251	64.1
Hispanic/Latino ^d	898	387	43.1	510	56.9
Native Hawaiian/Other Pacific Islander	2	1	50.2	1	49.8
White	2,398	923	38.5	1,475	61.5
Multiple races	125	53	42.5	72	57.5
Transmission category					
Male-to-male sexual contact	3,528	1,333	37.8	2,195	62.2
Injection drug use	554	214	38.7	340	61.3
Male-to-male sexual contact and injection drug use	163	58	35.8	104	64.2
Heterosexual contact ^e	1,188	492	41.4	696	58.6
Other ^f	17	8	45.0	9	55.0
Subtotal, males	5,450	2,106	38.6	3,344	61.4
Females					
Age at diagnosis (yr)					
50–54	876	281	32.1	595	67.9
55–59	631	233	36.9	398	63.1
60–64	324	127	39.3	196	60.7
65–69	130	49	38.0	81	62.0
70–74	70	24	34.6	45	65.4
75–79	29	11	36.9	18	63.1
80–84	8	3	36.8	5	63.2
≥85	2	1	46.6	1	53.4
Race/ethnicity					
American Indian/Alaska Native	16	5	33.5	11	66.5
Asian	20	6	31.0	14	69.0
Black/African American	1,304	412	31.6	893	68.4
Hispanic/Latino ^d	311	115	37.0	196	63.0
Native Hawaiian/Other Pacific Islander	4	3	75.1	1	24.9
White	359	158	44.0	201	56.0
Multiple races	55	30	54.0	25	46.0
Transmission category					
Injection drug use	320	118	36.8	202	63.2
Heterosexual contact ^e	1,743	609	35.0	1,134	65.0
Other ^f	7	3	37.6	4	62.4
Subtotal, females	2,070	730	35.2	1,341	64.8

Table 15a. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States (cont)

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
		Est. No. ^c	Est. No. ^c	%	Est. No. ^c
Both sexes					
Age at diagnosis (yr)					
50–54	3,235	1,191	36.8	2,045	63.2
55–59	2,184	823	37.7	1,361	62.3
60–64	1,170	442	37.7	729	62.3
65–69	549	233	42.4	316	57.6
70–74	237	91	38.6	146	61.4
75–79	93	40	43.1	53	56.9
80–84	37	13	35.7	24	64.3
≥85	14	2	14.8	12	85.2
Race/ethnicity					
American Indian/Alaska Native	32	12	37.0	20	63.0
Asian	80	41	51.6	39	48.4
Black/African American	3,255	1,112	34.2	2,144	65.8
Hispanic/Latino ^d	1,209	502	41.6	707	58.4
Native Hawaiian/Other Pacific Islander	6	4	66.8	2	33.2
White	2,758	1,081	39.2	1,676	60.8
Multiple races	180	83	46.0	97	54.0
Total^g	7,520	2,835	37.7	4,684	62.3

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Total column represents all persons with diagnosed HIV infection and is the sum of all persons with infection classified as Stage 3 (AIDS) or another stage.

^b Based on first CD4 test result or documentation of an AIDS-defining condition within 3 months after diagnosis of HIV infection.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15b. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
	Est. No. ^c	Est. No. ^c	%	Est. No. ^c	%
Males					
Age at diagnosis (yr)					
50–54	2,418	925	38.2	1,493	61.8
55–59	1,581	603	38.1	979	61.9
60–64	876	330	37.6	546	62.4
65–69	432	190	44.0	242	56.0
70–74	171	68	40.1	102	59.9
75–79	67	31	45.4	37	54.6
80–84	32	12	38.6	19	61.4
≥85	12	1	8.6	11	91.4
Race/ethnicity					
American Indian/Alaska Native	16	6	40.5	9	59.5
Asian	60	35	58.5	25	41.5
Black/African American	1,954	701	35.9	1,253	64.1
Hispanic/Latino ^d	1,035	440	42.5	594	57.5
Native Hawaiian/Other Pacific Islander	2	1	50.2	1	49.8
White	2,398	923	38.5	1,475	61.5
Multiple races	125	53	42.5	72	57.5
Transmission category					
Male-to-male sexual contact	3,562	1,345	37.7	2,218	62.3
Injection drug use	589	226	38.4	363	61.6
Male-to-male sexual contact and injection drug use	168	59	35.3	109	64.7
Heterosexual contact ^e	1,253	522	41.6	731	58.4
Other ^f	17	8	45.0	9	55.0
Subtotal, males	5,590	2,160	38.6	3,430	61.4
Females					
Age at diagnosis (yr)					
50–54	904	286	31.6	619	68.4
55–59	652	236	36.2	416	63.8
60–64	334	131	39.1	203	60.9
65–69	135	52	38.4	83	61.6
70–74	70	24	34.6	45	65.4
75–79	29	11	36.9	18	63.1
80–84	10	3	32.6	6	67.4
≥85	2	1	46.6	1	53.4
Race/ethnicity					
American Indian/Alaska Native	16	5	33.5	11	66.5
Asian	20	6	31.0	14	69.0
Black/African American	1,307	412	31.5	895	68.5
Hispanic/Latino ^d	373	129	34.5	244	65.5
Native Hawaiian/Other Pacific Islander	4	3	75.1	1	24.9
White	360	158	43.9	202	56.1
Multiple races	55	30	54.0	25	46.0
Transmission category					
Injection drug use	331	119	35.9	212	64.1
Heterosexual contact ^e	1,797	622	34.6	1,175	65.4
Other ^f	7	3	37.6	4	62.4
Subtotal, females	2,135	743	34.8	1,392	65.2

Table 15b. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas (cont)

	Total ^a	Stage 3 (AIDS) at diagnosis ^b			
		Yes		No	
		Est. No. ^c	%	Est. No. ^c	%
Both sexes					
Age at diagnosis (yr)					
50–54	3,323	1,211	36.4	2,112	63.6
55–59	2,234	839	37.6	1,394	62.4
60–64	1,210	460	38.1	749	61.9
65–69	567	242	42.7	325	57.3
70–74	240	93	38.5	148	61.5
75–79	96	41	42.8	55	57.2
80–84	41	15	37.2	26	62.8
≥85	14	2	14.8	12	85.2
Race/ethnicity					
American Indian/Alaska Native	32	12	37.0	20	63.0
Asian	80	41	51.6	39	48.4
Black/African American	3,261	1,113	34.1	2,148	65.9
Hispanic/Latino ^d	1,407	569	40.4	838	59.6
Native Hawaiian/Other Pacific Islander	6	4	66.8	2	33.2
White	2,759	1,081	39.2	1,678	60.8
Multiple races	180	83	46.0	97	54.0
Total^g	7,725	2,903	37.6	4,822	62.4

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Total column represents all persons with diagnosed HIV infection and is the sum of all persons with infection classified as Stage 3 (AIDS) or another stage.

^b Based on first CD4 test result or documentation of an AIDS-defining condition within 3 months after diagnosis of HIV infection.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 16a. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by age group and selected characteristics, 2013—United States

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
	Est. No. ^c	Est. No. ^c	%	Est. No. ^c	%
50–64 years of age at end of year					
Males					
Race/ethnicity					
American Indian/Alaska Native	13	4	34.1	8	65.9
Asian	54	32	59.4	22	40.6
Black/African American	1,695	595	35.1	1,100	64.9
Hispanic/Latino ^d	807	344	42.6	463	57.4
Native Hawaiian/Other Pacific Islander	2	1	50.2	1	49.8
White	2,082	793	38.1	1,288	61.9
Multiple races	107	45	42.2	62	57.8
Transmission category					
Male-to-male sexual contact	3,144	1,180	37.5	1,964	62.5
Injection drug use	488	182	37.4	306	62.6
Male-to-male sexual contact and injection drug use	146	51	34.9	95	65.1
Heterosexual contact ^e	973	399	41.0	574	59.0
Other ^f	7	2	33.6	5	66.4
Subtotal, males	4,759	1,815	38.1	2,944	61.9
Females					
Race/ethnicity					
American Indian/Alaska Native	14	3	22.6	11	77.4
Asian	17	4	24.4	13	75.6
Black/African American	1,154	367	31.8	788	68.2
Hispanic/Latino ^d	269	92	34.0	178	66.0
Native Hawaiian/Other Pacific Islander	4	3	75.1	1	24.9
White	324	144	44.5	180	55.5
Multiple races	49	29	58.9	20	41.1
Transmission category					
Injection drug use	291	104	35.9	186	64.1
Heterosexual contact ^e	1,537	536	34.8	1,001	65.2
Other ^f	3	1	38.6	2	61.4
Subtotal, females	1,831	641	35.0	1,190	65.0
Both sexes					
Race/ethnicity					
American Indian/Alaska Native	26	7	28.1	19	71.9
Asian	71	36	51.0	35	49.0
Black/African American	2,849	962	33.8	1,887	66.2
Hispanic/Latino ^d	1,076	435	40.5	641	59.5
Native Hawaiian/Other Pacific Islander	6	4	66.8	2	33.2
White	2,405	937	39.0	1,468	61.0
Multiple races	156	74	47.4	82	52.6
Subtotal, 50–64 years of age at end of year	6,590	2,456	37.3	4,134	62.7

Table 16a. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by age group and selected characteristics, 2013—United States (cont)

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
		Est. No. ^c	Est. No. ^c	%	Est. No. ^c
≥65 years of age at end of year					
Males					
Race/ethnicity					
American Indian/Alaska Native	3	2	66.6	1	33.4
Asian	6	3	51.0	3	49.0
Black/African American	256	105	41.0	151	59.0
Hispanic/Latino ^d	91	43	47.6	48	52.4
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0.0
White	317	130	41.0	187	59.0
Multiple races	17	8	44.4	10	55.6
Transmission category					
Male-to-male sexual contact	384	153	39.9	231	60.1
Injection drug use	65	32	48.5	34	51.5
Male-to-male sexual contact and injection drug use	16	7	44.3	9	55.7
Heterosexual contact ^e	215	94	43.5	121	56.5
Other ^f	10	5	53.3	5	46.7
Subtotal, males	691	291	42.2	400	57.8
Females					
Race/ethnicity					
American Indian/Alaska Native	2	2	100.0	0	0.0
Asian	3	2	66.8	1	33.2
Black/African American	150	45	30.1	105	69.9
Hispanic/Latino ^d	42	24	56.4	18	43.6
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0.0
White	36	14	40.1	21	59.9
Multiple races	6	1	16.9	5	83.1
Transmission category					
Injection drug use	30	13	45.5	16	54.5
Heterosexual contact ^e	206	74	35.7	133	64.3
Other ^f	3	1	36.6	2	63.4
Subtotal, females	239	88	36.9	151	63.1
Both sexes					
Race/ethnicity					
American Indian/Alaska Native	5	4	80.6	1	19.4
Asian	9	5	56.2	4	43.8
Black/African American	406	150	37.0	256	63.0
Hispanic/Latino ^d	133	67	50.4	66	49.6
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0.0
White	352	144	40.9	208	59.1
Multiple races	24	9	36.9	15	63.1
Subtotal, ≥65 years of age at end of year	930	380	40.8	551	59.2
Total^g	7,520	2,835	37.7	4,684	62.3

Reported numbers less than 12, as well as estimated numbers based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Total column represents all persons with diagnosed HIV infection and is the sum of all persons with infection classified as Stage 3 (AIDS) or another stage.

^b Based on first CD4 test result or documentation of an AIDS-defining condition within 3 months after diagnosis of HIV infection.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 16b. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
	Est. No. ^c	Est. No. ^c	%	Est. No. ^c	%
50–64 years of age at end of year					
Males					
Race/ethnicity					
American Indian/Alaska Native	13	4	34.1	8	65.9
Asian	54	32	59.4	22	40.6
Black/African American	1,698	596	35.1	1,102	64.9
Hispanic/Latino ^d	921	386	41.9	535	58.1
Native Hawaiian/Other Pacific Islander	2	1	50.2	1	49.8
White	2,082	793	38.1	1,288	61.9
Multiple races	107	45	42.2	62	57.8
Transmission category					
Male-to-male sexual contact	3,175	1,189	37.5	1,985	62.5
Injection drug use	519	193	37.2	325	62.8
Male-to-male sexual contact and injection drug use	152	52	34.3	100	65.7
Heterosexual contact ^e	1,024	421	41.1	603	58.9
Other ^f	7	2	33.6	5	66.4
Subtotal, males	4,876	1,858	38.1	3,018	61.9
Females					
Race/ethnicity					
American Indian/Alaska Native	14	3	22.6	11	77.4
Asian	17	4	24.4	13	75.6
Black/African American	1,157	367	31.7	790	68.3
Hispanic/Latino ^d	325	103	31.6	222	68.4
Native Hawaiian/Other Pacific Islander	4	3	75.1	1	24.9
White	325	144	44.3	181	55.7
Multiple races	49	29	58.9	20	41.1
Transmission category					
Injection drug use	301	105	35.1	195	64.9
Heterosexual contact ^e	1,586	546	34.4	1,041	65.6
Other ^f	3	1	38.6	2	61.4
Subtotal, females	1,890	652	34.5	1,238	65.5
Both sexes					
Race/ethnicity					
American Indian/Alaska Native	26	7	28.1	19	71.9
Asian	71	36	51.0	35	49.0
Black/African American	2,855	963	33.7	1,892	66.3
Hispanic/Latino ^d	1,246	489	39.2	757	60.8
Native Hawaiian/Other Pacific Islander	6	4	66.8	2	33.2
White	2,406	937	38.9	1,469	61.1
Multiple races	156	74	47.4	82	52.6
Subtotal, 50–64 years of age at end of year	6,766	2,510	37.1	4,256	62.9

Table 16b. Stage 3 (AIDS) at the time of diagnosis of HIV infection among adults aged 50 years and older, by sex and selected characteristics, 2013—United States and 6 dependent areas (cont)

	Stage 3 (AIDS) at diagnosis ^b				
	Total ^a	Yes		No	
		Est. No. ^c	Est. No. ^c	%	Est. No. ^c
≥65 years of age at end of year					
Males					
Race/ethnicity					
American Indian/Alaska Native	3	2	66.6	1	33.4
Asian	6	3	51.0	3	49.0
Black/African American	256	105	41.0	151	59.0
Hispanic/Latino ^d	114	54	47.7	60	52.3
Native Hawaiian/Other Pacific Islander	317	130	41.0	187	59.0
White	17	8	44.4	10	55.6
Multiple races					
Transmission category	388	156	40.1	232	59.9
Male-to-male sexual contact	70	33	46.9	37	53.1
Injection drug use	16	7	44.3	9	55.7
Male-to-male sexual contact and injection drug use	230	101	44.0	129	56.0
Heterosexual contact ^e	10	5	53.3	5	46.7
Other ^f	714	302	42.3	412	57.7
Subtotal, males					
Females					
Race/ethnicity					
American Indian/Alaska Native	2	2	100	0	0.0
Asian	3	2	66.8	1	33.2
Black/African American	150	45	30.1	105	69.9
Hispanic/Latino ^d	47	26	54.5	22	45.5
Native Hawaiian/Other Pacific Islander	36	14	40.1	21	59.9
White	6	1	16.9	5	83.1
Multiple races					
Transmission category					
Injection drug use	31	14	44.1	17	55.9
Heterosexual contact ^e	211	76	36.0	135	64.0
Other ^f	3	1	36.6	2	63.4
Subtotal, females					
Both sexes					
Race/ethnicity					
American Indian/Alaska Native	5	4	80.6	1	19.4
Asian	9	5	56.2	4	43.8
Black/African American	406	150	37.0	256	63.0
Hispanic/Latino ^d	162	80	49.7	81	50.3
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0.0
White	352	144	40.9	208	59.1
Multiple races	24	9	36.9	15	63.1
Subtotal, ≥65 years of age at end of year					
Total^g	7,725	2,903	37.6	4,822	62.4

Reported numbers less than 12, as well as estimated numbers based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Total column represents all persons with diagnosed HIV infection and is the sum of all persons with infection classified as Stage 3 (AIDS) or another stage.

^b Based on first CD4 test result or documentation of an AIDS-defining condition within 3 months after diagnosis of HIV infection.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.